

EXISTENCE OF QUASINORMAL MODES FOR KERR–ADS BLACK HOLES

ORAN GANNOT

ABSTRACT. This paper establishes the existence of quasinormal frequencies converging exponentially to the real axis for the Klein–Gordon equation on a Kerr–AdS spacetime when Dirichlet boundary conditions are imposed at the conformal boundary. The proof is adapted from results in Euclidean scattering about the existence of scattering poles generated by sequences of time-periodic approximate solutions to the wave equation.

1. INTRODUCTION

Recent years have seen substantial progress in the analysis of linear fields on asymptotically anti-de Sitter (aAdS) backgrounds. Understanding the boundedness and decay of solutions to the linear wave equation is a prerequisite for studying nonlinear (in)stability of aAdS backgrounds. Furthermore, linear fields play a distinguished role in the AdS/CFT correspondence.

One common way of describing linear perturbations of black holes is through the quasinormal frequency (QNF) spectrum, consisting of complex frequencies λ associated with oscillating and decaying quasinormal mode (QNM) solutions to the wave equation. Despite a sizable literature concerning the QNFs of aAdS black holes, there are still many open questions in the mathematical study of these objects.

One notable conjecture is that aAdS black holes should display QNFs rapidly converging to the real axis. This phenomenon was first observed for the Schwarzschild–AdS solution through numerical and formal WKB analysis [8, 12]. The existence of such highly damped modes is consistent with at most logarithmic local energy decay in time for solutions of the wave equation [14, 18, 19]. In addition, the asymptotic relation between these QNFs and the spectrum of global AdS provides a link between the conjectured nonlinear instability of global AdS and that of Kerr–AdS [1, 2, 3, 4, 5, 7, 6, 8, 9, 18].

In [14], the existence of QNFs converging exponentially to the real axis was rigorously established for Schwarzschild–AdS black holes. This was based on a construction of quasimodes (not to be confused with QNMs), which are time-periodic approximate solutions to the Klein–Gordon equation. Due to the spherical symmetry of the

Schwarzschild–AdS metric, the wave equation separates into a family of one dimensional equations indexed by angular momenta ℓ , each of which fits into the framework of classical scattering theory on the half-line. The quasimodes constructed in [14] reflect the existence of null-geodesics which are trapped between the conformal boundary and an effective potential barrier. By applying general results of Tang–Zworski [29] from Euclidean scattering, it was possible to conclude the existence of QNFs converging exponentially to the real axis as $\ell \rightarrow \infty$.

For the Kerr–AdS solution, it is more difficult to demonstrate the existence of long-lived QNMs. Due to the more complicated structure of the separated equations, a WKB analysis is harder to perform. Furthermore, the author is not aware of any numerical studies of QNFs for Kerr–AdS in the high frequency limit.

Nevertheless exponentially accurate quasimodes have been constructed for Kerr–AdS metrics by Holzegel–Smulevici [19]; their motivation was to establish a logarithmic lower bound for energy decay. At a linearized level, the construction of [19] is the same as in [14]. A natural question is whether the methods of Tang–Zworski can be adapted to deduce the existence of QNFs converging to the real axis from these quasimodes.

Theorem 1 (Main Theorem). *Fix a cosmological constant $\Lambda < 0$, black hole mass $M > 0$, rotation speed $a \in \mathbb{R}$ satisfying $|a|^2 < 3/|\Lambda|$, and Klein–Gordon mass $\nu > 0$. Let $X_\delta = (r_+ - \delta, \infty) \times \mathbb{S}^2$ for $\delta > 0$ sufficiently small, and let t^* be the Kerr-star time coordinate. Then there exists a sequence of complex numbers and smooth functions*

$$\lambda_\ell \in \mathbb{C}, \quad u_\ell \in C^\infty(X_\delta), \quad \ell \geq L$$

for some $L \geq 0$ with the following properties.

- (1) The functions $v_\ell := e^{-i\lambda_\ell t^*} u_\ell$ solve the Klein–Gordon equation

$$\square_g v_\ell + \frac{|\Lambda|}{3}(\nu^2 - 9/4)v_\ell = 0.$$

- (2) The complex frequencies λ_ℓ satisfy

$$\ell/C < \operatorname{Re} \lambda_\ell < C\ell, \quad 0 < -\operatorname{Im} \lambda_\ell < e^{-\ell/D}$$

for some $C, D > 0$.

- (3) Each u_ℓ is smooth up to $\{r = r_+ - \delta\}$, and the restriction of u_ℓ to $X_0 = (r_+, \infty) \times \mathbb{S}^2$ is nonzero.

- (4) Each u_ℓ satisfies

$$\int_{X_\delta} |u_\ell|^2 r^{-1} dS_t < \infty,$$

where dS_t is the surface measure induced on X_δ by g , and moreover

$$\lim_{r \rightarrow \infty} r^{3/2-\nu} u_\ell = 0.$$

- (5) Each u_ℓ is axisymmetric in the sense that $\partial_\phi u = 0$, where ϕ is the azimuthal angle on $\mathbb{S}^2$.

The frequencies λ_ℓ in Theorem 1 are QNFs, and u_ℓ are associated QNMs. The theorem is deduced from the existence of *real* frequencies $\lambda_\ell^\# \in \mathbb{R}$ and nonzero functions $u_\ell^\# \in C_c^\infty(X_0)$ for which (2), (4), (5) hold, such that (1) is approximately satisfied (see Theorem 3 below for a more precise statement regarding these quasimodes).

Remark. (1) The functions v_ℓ are smooth solutions to the Klein–Gordon equation in a region extending past the event horizon. This reflects the outgoing nature of QNMs. Since u_ℓ does not vanish outside the event horizon one also obtains a nonzero solution to the Klein–Gordon equation by restricting to the black hole exterior.

(2) The square integrability condition (4) constrains the growth of u_ℓ as $r \rightarrow \infty$. In fact, u_ℓ has a conormal asymptotic expansion near the conformal boundary determined by the indicial roots of the Klein–Gordon operator [15, Proposition 4.17]. The coefficient of $r^{\nu-3/2}$ vanishes, which is a type of Dirichlet boundary condition in this setting.

As will be clear from the proof, Theorem 1 is a *black box* in the sense that any sequence of quasimodes satisfying the conditions of Theorem 3 can be plugged into the machinery to obtain a corresponding sequence of QNFs. Furthermore, there is a relationship

$$|\lambda_\ell - \lambda_\ell^\#| \leq e^{-\ell/C} \quad (1.1)$$

for some $C > 0$. Thus any description of $\lambda_\ell^\#$ modulo $\mathcal{O}(\ell^{-\infty})$ gives a corresponding description for $\text{Re } \lambda_\ell$ as $\ell \rightarrow \infty$. The imprecise localization of $\text{Re } \lambda_\ell$ in Theorem 1 is therefore only due to the inexact nature of the quasimodes constructed in [19]; this should be compared to the main theorem of [14] in the simpler Schwarzschild–AdS setting, where $\text{Re } \lambda_\ell$ admits an asymptotic expansion in powers of $\ell^{-1/2}$.

Remark. (1) Exponential accuracy of the quasimodes is not necessary to deduce the existence of QNFs — see the proof of Theorem 1, as well as [27, 28, 29] for more general results in the Euclidean setting. Less accurate quasimodes could potentially result in slower convergence to the real axis, as well as a weaker version of (1.1).

(2) Theorem 1 still applies if the quasimodes are supported on finitely many eigenspaces of D_ϕ (uniformly in ℓ).

(3) Quasimodes satisfying more general self-adjoint boundary conditions also yield a version of Theorem 1 — see the discussion in Section 2.2, as well as the statements of Propositions 2.1, 2.2, 2.3.

A natural question is to what extent the passage from quasimodes to QNFs depends on the exact form of the Kerr–AdS metric. Observe already that axial symmetry of the

Kerr–AdS metric plays an important role in the statement of Theorem 1. This allows one to compensate for the fact that the Killing field ∂_t is not timelike near the event horizon for $a \neq 0$; see Propositions 2.1, 2.2 below. For the full range of parameters $|a|^2 < 3/|\Lambda|$, these propositions apply to stationary, *axisymmetric* perturbations of the metric (throughout, perturbations are assumed to be small).

On the other hand, the final ingredient in the proof of Theorem 1, Proposition 2.3 below, is always stable under stationary perturbations of the metric. The analysis is based on a general microlocal framework developed by Vasy [30], which is highly robust — see [30, Section 2.7] for a precise discussion.

2. PROOF OF THEOREM 1

The proof of Theorem 1 relies on three key results, Propositions 2.1, 2.2, 2.3, stated in the next section. The proof of Proposition 2.3 is delayed until Section 3, while Proposition 2.1 is proved in Section 4 at the end of the paper.

2.1. Kerr–AdS metric. The Kerr–AdS metric is determined by three parameters (Λ, M, a) , where $\Lambda < 0$ is the negative cosmological constant, $M > 0$ is the black hole mass, and $a \in \mathbb{R}$ is the angular momentum per unit mass. It is always possible to choose units such that $\Lambda = -3$, in which case the rotation speed is required to satisfy the regularity condition $|a| < 1$. Introduce the quantities

$$\begin{aligned}\Delta_r &= (r^2 + a^2)(1 + r^2) - 2Mr; & \Delta_\theta &= 1 - a^2 \cos^2 \theta; \\ \varrho^2 &= r^2 + a^2 \cos^2 \theta; & \alpha &= a^2.\end{aligned}$$

When it exists, the largest positive root of Δ_r is denoted by r_+ . The Kerr–AdS metric is given in Boyer–Lindquist coordinates by the expression

$$\begin{aligned}g &= -\varrho^2 \left(\frac{dr^2}{\Delta_r} + \frac{d\theta^2}{\Delta_\theta} \right) - \frac{\Delta_\theta \sin^2 \theta}{\varrho^2(1 - \alpha)^2} (a dt - (r^2 + a^2) d\phi)^2 \\ &\quad + \frac{\Delta_r}{\varrho^2(1 - \alpha)^2} (dt - a \sin^2 \theta d\phi)^2.\end{aligned}$$

Here $(\theta, \phi) \in (0, \pi) \times \mathbb{R}/(2\pi\mathbb{Z})$ are spherical coordinates on $\mathbb{S}^2$, while $t \in \mathbb{R}$ and $r \in (r_+, \infty)$. The dual metric g^{-1} is given by

$$\begin{aligned}g^{-1} &= \frac{\Delta_r}{\varrho^2} D_r^2 + \frac{\Delta_\theta}{\varrho^2} D_\theta^2 + \frac{(1 - \alpha)^2}{\varrho^2 \Delta_\theta \sin^2 \theta} (a \sin^2 \theta D_t + D_\phi)^2 \\ &\quad - \frac{(1 - \alpha)^2}{\varrho^2 \Delta_r} ((r^2 + a^2) D_t + a D_\phi)^2.\end{aligned}$$

The metric g becomes singular at the event horizon $\{r = r_+\}$, but this can be remedied by a change of variables. Set

$$t^\star = t + F_t(r); \quad \phi^\star = \phi + F_\phi(r), \tag{2.1}$$

where F_t, F_ϕ are smooth functions on (r_+, ∞) such that

$$\partial_r F_t(r) = \frac{1-\alpha}{\Delta_r}(r^2 + a^2) + f_+(r), \quad \partial_r F_\phi(r) = a \frac{1-\alpha}{\Delta_r}. \quad (2.2)$$

Here f_+ is an arbitrary smooth function, and F_t, F_ϕ are chosen to vanish at infinity. The dual metric in (t^*, r, θ, ϕ^*) coordinates reads

$$\begin{aligned} \varrho^2 g^{-1} = & -\Delta_r (\partial_r + f_+ \partial_{t^*})^2 - \Delta_\theta \partial_\theta^2 - 2(1-\alpha) (\partial_r + f_+ \partial_{t^*}) ((r^2 + a^2) \partial_{t^*} + a \partial_{\phi^*}) \\ & - \frac{(1-\alpha)^2}{\Delta_\theta \sin^2 \theta} (a \sin^2 \theta \partial_{t^*} + \partial_{\phi^*})^2, \end{aligned} \quad (2.3)$$

The identification $\partial_t = \partial_{t^*}, \partial_\phi = \partial_{\phi^*}$ under $(t, r, \theta, \phi) \mapsto (t^*, r, \theta, \phi^*)$ will be used throughout. Furthermore, write

$$T = \partial_t, \quad \Phi = \partial_\phi.$$

Given $\delta \geq 0$ set $X_\delta = (r_+ - \delta, \infty) \times \mathbb{S}^2$ and $\mathcal{M}_\delta = X_\delta \times \mathbb{R}_{t^*}$. Then for sufficiently small $\delta > 0$ the metric g admits an extension from $\mathcal{M}_0$ to $\mathcal{M}_\delta$. The function f_+ is chosen such that dt^* is timelike and $f_+ = \mathcal{O}(r^{-2})$ as $r \rightarrow \infty$. One explicit choice is $f_+(r) = (\alpha - 1)/(r^2 + 1)$, but any choice of f_+ with these properties is just as good.

Finally, it must also be assumed that the root of Δ_r at $r = r_+$ is simple, or equivalently $\partial_r \Delta_r(r_+) > 0$. This implies that the surface gravity κ of the event horizon is positive. Explicitly

$$\kappa = \frac{\partial_r \Delta_r(r_+)}{2(1-\alpha)(r_+^2 + a^2)}, \quad (2.4)$$

which is manifestly positive.

2.2. Quasinormal modes. To define the stationary Klein–Gordon operator, begin by identifying X_δ with the time slice $\{t^* = 0\} \subseteq \mathcal{M}_\delta$. Let $P(\lambda)$ denote the operator obtained from $\square_g + \nu^2 - 9/4$ by replacing D_t with $-\lambda \in \mathbb{C}$. Because the metric is stationary, $P(\lambda)$ is a well defined second order operator on X_δ . Given $u \in C^\infty(X_\delta)$ and $\lambda \in \mathbb{C}$,

$$P(\lambda)u = e^{i\lambda t^*} (\square_g + \nu^2 - 9/4) e^{-i\lambda t^*} u.$$

Furthermore, given $k \in \mathbb{Z}$ let $P(\lambda, k)$ denote the operator obtained from $P(\lambda)$ by replacing D_ϕ with k . Since D_ϕ is also Killing, $P(\lambda, k)$ preserves the space of distributions

$$\mathcal{D}'_k(X_\delta) = \{u \in \mathcal{D}'(X_\delta) : (\Phi - ik)u = 0\}.$$

Define $\mathcal{L}^2(X_\delta)$ as the space of distributions u for which

$$\|u\|_{\mathcal{L}^2(X_\delta)} = \int_{X_\delta} |u|^2 r^{-1} dS_t < \infty,$$

where dS_t is the surface measure induced on X_δ by g . The energy space is defined with respect to the conjugated differential (on X_δ),

$$\tilde{d}u = r^{\nu-3/2}d(r^{3/2-\nu}u).$$

Thus $\mathcal{H}^1(X_\delta)$ is defined as the space of distributions u for which

$$\|u\|_{\mathcal{H}^1(X_\delta)} = \int_{X_\delta} \left(|u|^2 + r^2 g^{-1}(\tilde{d}u, \tilde{d}\bar{u}) \right) r^{-1} dS_t < \infty,$$

observing that the restriction of g to the tangent bundle of X_δ is positive definite. The various powers of r appearing as weights are motivated by natural energy identities [17, 18, 21, 31, 32], see also Section 3.4 of this paper.

If $\nu \in (0, 1)$, then boundary conditions must be imposed at the conformal boundary $X_\delta \cap \mathcal{J} = \{r^{-1} = 0\}$ to obtain the Fredholm property for $P(\lambda)$. Given $u \in \mathcal{L}^2(X_\delta)$ such that $P(\lambda)u \in r^{-2}\mathcal{L}^2(X_\delta)$, the following boundary values are well defined:

$$\gamma_- u = \lim_{r \rightarrow \infty} r^{3/2-\nu}u, \quad \gamma_+ u = \lim_{r \rightarrow \infty} r^{2\nu+1}\partial_r(r^{3/2-\nu}u). \quad (2.5)$$

Thus $\gamma_- u$ and $\gamma_+ u$ are analogues of Dirichlet and Neumann data for u . This can be proved by an analogue of the usual partial hypoellipticity at the boundary argument [22, Theorem B.2.9]. In fact, $P(\lambda)$ is elliptic at the boundary $X_\delta \cap \mathcal{J}$ in the sense of Bessel operators [15, Section 1.5], so from the remark following [15, Lemma 4.13] and [15, Proposition 3.6] one has the sharper Sobolev regularity $\gamma_\pm u \in H^{\mp\nu}(\mathbb{S}^2)$ [15, Section 3.4].

Throughout this paper only self-adjoint Dirichlet or Robin boundary conditions are considered. The trace operator B is therefore

$$B = \gamma_- \text{ or } B = \gamma_+ + \beta\gamma_-,$$

where $\beta \in C^\infty(X_\delta \cap \mathcal{J}; \mathbb{R})$ is a real valued on the conformal boundary of X_δ . Then define the domain

$$\mathcal{X} = \begin{cases} \{u \in \mathcal{H}^1(X_\delta) : P(0)u \in r^{-2}\mathcal{L}^2(X_\delta)\} & \text{if } \nu \geq 1, \\ \{u \in \mathcal{H}^1(X_\delta) : P(0)u \in r^{-2}\mathcal{L}^2(X_\delta) \text{ and } Bu = 0\} & \text{if } \nu \in (0, 1). \end{cases} \quad (2.6)$$

This is a Hilbert space for the norm $\|u\|_{\mathcal{X}} = \|u\|_{\mathcal{H}^1(X_\delta)} + \|P(0)u\|_{r^{-2}\mathcal{L}^2(X_\delta)}$.

QNFs are defined in the following theorem, which is proved in [13] and also [32] under slightly different hypotheses.

Theorem 2. *For each $\nu > 0$ the operator $P(\lambda) : \mathcal{X} \rightarrow r^{-2}\mathcal{L}^2(X_\delta)$ is Fredholm of index zero in the half-plane $\{\text{Im } \lambda > -\frac{1}{2}\kappa\}$, where the surface gravity $\kappa > 0$ is given by (2.4). Furthermore,*

$$R(\lambda) := P(\lambda)^{-1} : r^{-2}\mathcal{L}^2(X_\delta) \rightarrow \mathcal{X}$$

is a meromorphic family of operators in $\{\text{Im } \lambda > -\frac{1}{2}\kappa\}$, which is holomorphic in any angular sector of the upper half-plane provided $|\lambda|$ is sufficiently large.

QNFs are defined as poles of the meromorphic family $R(\lambda)$. More information is needed about possible QNFs in the upper half-plane — this is closely related to the boundedness of solutions to the Klein–Gordon equation. One remarkable property of rotating Kerr–AdS metrics is that for $|a| < r_+^2$ there exists a timelike Killing field K on $\mathcal{M}_0$ which is null on the horizon:

$$K = T + \Omega\Phi, \quad \Omega = \frac{a}{r_+^2 + a^2}.$$

The existence of such a vector field eliminates possible superradiant phenomena. For black holes satisfying the Hawking–Reall bound $|a| < r_+^2$, boundedness [20] (and in fact logarithmic decay [18]) is known for solutions to the Klein–Gordon under Dirichlet boundary conditions. If the condition $|a| < r_+^2$ is violated, then it is possible to construct mode solutions $e^{-i\lambda t^*}u$ which grow exponentially in time, namely $\text{Im } \lambda > 0$ [10].

Remark. *Interestingly, even for Neumann boundary conditions boundedness has not been established for the expected range of black hole parameters $|a| < r_+^2$, see the conjecture in [20, Section 5].*

For mode solutions $e^{-i\lambda t^*}u$ many of the delicate issues involving lower order terms and boundary conditions are overcome in the high frequency limit. In fact by working at a fixed axial mode it is not even necessary to restrict below the Hawking–Reall bound; this is of course only possible because of the axisymmetry of the Kerr–AdS metric. In that case the Robin function β should satisfy $\Phi\beta = 0$ as well.

Note that $R(\lambda)$ decomposes as direct sum of operators

$$R(\lambda, k) : r^{-2}\mathcal{L}^2(X_\delta) \cap \mathcal{D}'_k(X_\delta) \rightarrow \mathcal{X} \cap \mathcal{D}'_k(X_\delta),$$

where $R(\lambda, k)$ is just the restriction of $R(\lambda)$ to $r^{-2}\mathcal{L}^2(X_\delta) \cap \mathcal{D}'_k(X_\delta)$. In particular, λ_0 is a QNF if and only if there exists $k_0 \in \mathbb{Z}$ such that λ_0 is a pole of $R(\lambda, k_0)$. The following crucial proposition quantifies the absence of QNFs at a fixed axial mode in the upper half-plane at high frequencies.

Proposition 2.1. *Given $k \in \mathbb{Z}$ there exists $C_0 > 0$ such that if $\lambda \in (C_0, \infty) + i(0, \infty)$, then*

$$\|u\|_{\mathcal{L}^2(X_0)} \leq \frac{C}{|\lambda| \text{Im } \lambda} \|P(\lambda, k)u\|_{r^{-2}\mathcal{L}^2(X_0)} \quad (2.7)$$

for each $u \in \mathcal{H}^1(X_0) \cap \mathcal{D}'_k(X_0)$ such that: i) $P(0)u \in r^{-2}\mathcal{L}^2(X_0)$, and ii) if $\nu \in (0, 1)$, then $Bu = 0$.

Proposition 2.1 is stated for functions on the exterior time slice X_0 rather than the extended region X_δ . On the other hand, [13, Theorem 3] combined with Proposition

[2.1](#) implies that $R(\lambda, k)$ has no poles in the region $\lambda \in (C_0, \infty) + i(0, \infty)$. Moreover, if $\mathcal{R} : \mathcal{L}^2(X_\delta) \rightarrow \mathcal{L}^2(X_0)$ is the restriction operator, then [\(2.7\)](#) implies

$$\|\mathcal{R}R(\lambda, k)f\|_{\mathcal{L}^2(X_0)} \leq \frac{C}{|\lambda| \operatorname{Im} \lambda} \|f\|_{r^{-2}\mathcal{L}^2(X_\delta)} \quad (2.8)$$

for each $f \in r^{-2}\mathcal{L}^2(X_\delta)$ and $\lambda \in (C_0, \infty) + i(0, \infty)$. The constant $C > 0$ in [\(2.8\)](#) a priori may depend on $k \in \mathbb{Z}$.

Remark. *Proposition [2.1](#) is also valid without restricting to a fixed axial mode provided the Hawking–Reall bound holds (as will be evident from the proof).*

It is also important to know that there are no QNFs on the real axis. The proof also exploits axisymmetry of the Kerr–AdS metric, hence the Robin function should satisfy $\Phi\beta = 0$ as well.

Proposition 2.2. *Let B be as in Proposition [2.1](#). Suppose that*

$$u \in \mathcal{H}^1(X_0) \cap C^\infty(X_0)$$

satisfies: i) u is smooth up to $\{r = r_+\}$, ii) $P(\lambda_0)u = 0$, iii) if $\nu \in (0, 1)$, then $Bu = 0$. If $\lambda_0 \in \mathbb{R} \setminus 0$ then $u = 0$.

Proof. According to [\[32, Lemma A.1\]](#) the solution u vanishes in a neighborhood of $\{r = r_+\}$ within X_0 . As in that lemma, one would like to apply some type of unique continuation result to conclude that once away from the horizon, u must vanish everywhere. This is known to be a difficult problem in view of possible trapping within the ergoregion where $P(\lambda)$ fails to be elliptic [\[23\]](#). To work around this, define the Riemannian metric

$$h = \varrho^2 \left(\frac{1}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sin^2 \theta}{(1 - \alpha)^2} d\phi^2 \right),$$

and observe that the difference between $P(\lambda)$ and the operator $\varrho^{-2}\Delta_r D_r^2 - \Delta_h$ is of first order modulo the second order term $2a\varrho^{-2}(1 - \alpha)D_r D_\phi$. Set

$$\tilde{P}(\lambda, k) = \varrho^{-2}\Delta_r D_r^2 - \Delta_h + e^{-ik\phi^*} (P(\lambda) - (\varrho^{-2}\Delta_r D_r^2 - \Delta_h)) e^{ik\phi^*}.$$

Thus $\tilde{P}(\lambda, k)$ is elliptic on $\{r > r_+\}$, and furthermore

$$\tilde{P}(\lambda, k)u = P(\lambda, k)u \quad (2.9)$$

for each $u \in \mathcal{D}'_k(X_0)$.

Now decompose u as an $\mathcal{L}^2(X_0)$ orthogonal sum $u = \sum_{k \in \mathbb{Z}} u_k$, where $u_k \in \mathcal{H}^1(X_0) \cap C^\infty(X_0) \cap \mathcal{D}'_k(X_0)$ is smooth up to $\{r = r_+\}$, satisfies $P(\lambda_0, k)u_k = 0$, and $Bu_k = 0$ if $\nu \in (0, 1)$. Unique continuation holds for $\tilde{P}(\lambda, k)$, hence each u_k vanishes by [\(2.9\)](#) since it satisfies $\tilde{P}(\lambda, k)u_k = 0$ and u_k vanishes in a neighborhood of the horizon. $\square$

Again referring to [13, Theorem 3], the previous proposition implies that $R(\lambda)$ does not have any poles for $\lambda \in \mathbb{R} \setminus 0$.

The crucial ingredient used to prove Theorem 1 is an exponential bound on $R(\lambda)$ in a strip away from suitable neighborhoods of the poles of $R(\lambda)$. At this stage it is convenient to introduce the semiclassical rescaling. Given a parameter $h > 0$, set $z = h\lambda$ and

$$P_h(z) = h^2 P(h^{-1}z), \quad R_h(z) = P_h(z)^{-1},$$

and similarly for $P_h(z, k)$ and $R_h(z, k)$. For the next proposition the Robin function β is not required to satisfy $\Phi\beta = 0$; in fact, no integrability properties of the metric are used at all. Fix intervals $[a, b] \subseteq (0, \infty)$ and $[C_-, C_+] \subseteq (-\kappa/2, \infty)$. Given $\varepsilon > 0$, define

$$\Omega_\varepsilon(h) = [a - \varepsilon h, b + \varepsilon h] + ih[C_- + \varepsilon, C_+ + \varepsilon]. \quad (2.10)$$

Also write $\Omega(h) := \Omega_0(h)$.

Proposition 2.3. *Fix $\varepsilon > 0$ and let $\{z_j\}$ denote the poles of $R_h(z)$ in $\Omega_\varepsilon(h)$. Then there exists $A > 0$ such that given any function $0 < S(h) = o(h)$,*

$$\|R_h(z)\|_{\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{L}^2(X_\delta)} < \exp(Ah^{-12} \log(1/S(h))), \quad (2.11)$$

for

$$z \in \Omega(h) \setminus \bigcup_j B(z_j, S(h))$$

and h sufficiently small.

Proposition 2.3 is proved in Section 3.6. Finally, the quasimode construction of [19] is reviewed:

Theorem 3 ([19, Theorem 1.2]). *There exists a sequence $\lambda_\ell^\sharp \in \mathbb{R}$ and*

$$u_\ell^\sharp \in \mathcal{H}^1(X_\delta) \cap C^\infty(X_\delta) \cap \mathcal{D}'_0(X_\delta)$$

with the following properties.

- (1) *There exists $C > 0$ such that $\ell/C < \lambda_\ell^\sharp < C\ell$.*
- (2) *There exists $[r_1, r_2] \subseteq (r_+, \infty)$ such that $\text{supp } u_\ell^\sharp \subseteq (r_1, \infty)$ and $\text{supp } P(\lambda_\ell^\sharp, 0)u_\ell^\sharp \subseteq (r_1, r_2)$.*
- (3) *The functions u_ℓ are normalized exponentially accurate quasimodes in the sense that*

$$\|u_\ell\|_{\mathcal{L}^2(X_\delta)} = 1, \quad \|P(\lambda_\ell^\sharp, 0)u_\ell\|_{r^{-2}\mathcal{L}^2(X_\delta)} \leq e^{-\ell/C}$$

for some $C > 0$. Furthermore, if $\nu \in (0, 1)$, then $\gamma_- u_\ell = 0$.

As remarked in [19, Footnote 8], the boundary condition $B = \gamma_-$ in [19, Theorem 1.2] could also be replaced by a Robin boundary condition of the form $B = \gamma_+ + \beta\gamma_-$, where $\beta \in \mathbb{R}$ is a real constant. The only difference is that the Hardy inequality used in [19] now holds modulo a boundary term which is negligible in the semiclassical limit (after applying a trace identity [32, Lemma B.1]).

The proof of Theorem 1 can now be finished by the same arguments as in the work of Tang–Zworski [29] with refinements by Stefanov [27, 28].

Proof of Theorem 1. Define a semiclassical parameter h by $h^{-1} := \lambda_\ell^\#$ and set $u(h) := u_\ell$. Then, $h \rightarrow 0$ as $\ell \rightarrow \infty$.

Suppose that $R_h(z, 0)$ was holomorphic in the rectangle

$$[1 - 2w(h) - S(h), 1 + 2w(h) + S(h)] + i[-Ah^{-12} \log(1/S(h))S(h), S(h)], \quad (2.12)$$

where $A > 0$ is as in Proposition 2.3, and $w(h)$, $S(h)$ are to be specified. Then

$$F(z) := \mathcal{R}R_h(z, 0) : r^{-2}\mathcal{L}^2(X_\delta) \rightarrow \mathcal{L}^2(X_0)$$

is holomorphic in the smaller rectangle

$$\Sigma(h) = [1 - 2w(h), 1 + 2w(h)] + i[-Ah^{-12} \log(1/S(h))S(h), S(h)],$$

and satisfies the operator norm estimates

$$\|F(z)\| < \begin{cases} C/\operatorname{Im} z & \text{for } z \in \Sigma(h) \cap \{\operatorname{Im} z > 0\}, \\ e^{Ah^{-12} \log(1/S(h))} & \text{for } z \in \Sigma(h). \end{cases}$$

from Propositions 2.1, 2.3. Applying the semiclassical maximum principle [28, Lemma 1], it follows that

$$\|F(z)\| < e^3/S(h) \text{ for } z \in [1 - w(h), 1 + w(h)]$$

provided $w(h)$, $S(h)$ are chosen so that $e^{-B/h} < S(h) < 1$ for some $B > 0$ and

$$2Aph^{-12} \log(1/h) \log(1/S(h))S(h) \leq w(h). \quad (2.13)$$

Define $\varepsilon(h) = \|P_h(1, 0)u(h)\|_{L^2(X_\delta)}$. Then,

$$1 = \|u(h)\|_{\mathcal{L}^2(X_\delta)} = \|F(1)P_h(1, 0)u(h)\|_{\mathcal{L}^2(X_0)} < e^3\varepsilon(h)/S(h).$$

This is a contradiction if $S(h) = 2e^3\varepsilon(h)$ for example. Thus there must exist a pole in the rectangle (2.12), which furthermore must lie in $\{\operatorname{Im} z < 0\}$ because of Propositions 2.1, 2.2. Defining $w(h)$ by the left hand side of (2.13), it follows that $R_h(z, 0)$ has a pole $z(h)$ such that

$$|z(h) - 1| < Ch^{-13} \log(1/h)\varepsilon(h), \quad 0 < -\operatorname{Im} z(h) < Ch^{-13}\varepsilon(h).$$

This is an even stronger statement than that of Theorem 1. □

The rest of the paper is dedicated to the proving Propositions 2.1 and 2.3.

3. EXPONENTIAL BOUNDS ON THE RESOLVENT

The first goal is to prove Proposition 2.3. For this an approximate inverse is constructed for $P(\lambda)$ modulo an error of Schatten class. The approximate inverse is built up from local parametrices which invert $P(\lambda)$ near the event horizon and near the conformal boundary. This is similar to the black-box approach of Sjöstrand–Zworski [26].

3.1. Microlocal analysis of the stationary operator. In order to construct a local parametrix in a neighborhood of the event horizon using methods of Vasy [30], one needs to understand the Hamilton flow of the symbol of $P_h(z)$ near its characteristic set.

It is convenient to view the (rescaled) flow on a compactified phase space $\overline{T}^*X_\delta$. The fibers of $\overline{T}^*X_\delta$ are obtained by gluing a sphere at infinity to the fibers of T^*X_δ , so $\overline{T}^*X_\delta$ is a disk bundle whose interior is identified with T^*X_δ — see [25, 30] as well as [11, Appendix E] for more details. If $|\cdot|$ is a smooth norm on the fibers of T^*X_δ , then a function on $\overline{T}^*X_\delta$ is smooth in a neighborhood of $\partial\overline{T}^*X_\delta$ if it is smooth in the polar coordinates $(x, \rho = |\xi|^{-1}, \omega = |\xi|^{-1}\xi)$, where $(x, \xi) \in T^*X_\delta$.

Throughout, $\text{Im } z$ will satisfy $|\text{Im } z| < Ch$ for some $C > 0$. In the semiclassical regime one may therefore assume that z is in fact real. The semiclassical principal symbol $p = \sigma_h(P_h(z))$ is given by

$$p(x, \xi; z) = -g^{-1}(\xi \cdot dx - z dt^*, \xi \cdot dx - z dt^*),$$

where $\xi \cdot dx$ is the canonical one-form on X_δ .

Note that $\langle \xi \rangle^{-2} p$ extends smoothly to a function on $\overline{T}^*X_\delta$ — here $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$ with respect to the fixed norm $|\cdot|$ on T^*X_δ . Furthermore, the rescaled Hamilton vector field $\langle \xi \rangle^{-1} H_p$ extends to a smooth vector field on $\overline{T}^*X_\delta$ which is tangent to $\partial\overline{T}^*X_\delta$. The Hamilton flow of p will refer to integral curves of $\langle \xi \rangle^{-1} H_p$ in this compactified picture.

The characteristic set of p is given by $\Sigma = \{\langle \xi \rangle^{-2} p = 0\} \subseteq \overline{T}^*X_\delta$, and its complement is the elliptic set. Since $X_\delta \subseteq \mathcal{M}_\delta$ is spacelike, it is a general fact about Lorentzian metrics that for each $z \in \mathbb{R} \setminus 0$ the characteristic set is the union of two disjoint sets $\Sigma = \Sigma_\pm$, where

$$\Sigma_\pm = \Sigma \cap \{\pm \langle \xi \rangle^{-1} g(\xi \cdot dx - z dt^*, dt^*) < 0\}$$

are the backwards and forwards light cones [30, Section 3.2]. Each of these sets is invariant under the Hamilton flow. Finally, let

$$\widehat{\Sigma} = \Sigma \cap \partial\overline{T}^*X_\delta, \quad \widehat{\Sigma}_\pm = \Sigma_\pm \cap \partial\overline{T}^*X_\delta.$$

If $\partial\overline{T}^*X_\delta$ is identified with the unit sphere bundle S^*X_δ via a norm on the fibers, then $\widehat{\Sigma}$ corresponds to the characteristic set of the homogeneous principal symbol of $P(\lambda)$ (as a non-semiclassical differential operator) within S^*X .

If $x \in X_\delta$ and the vector field T is timelike at x , then p is elliptic near the fiber $\partial\overline{T}_x^*X_\delta$. Therefore the projection of $\widehat{\Sigma}$ onto the base space X_δ is contained within the ergoregion, namely the set where T is not timelike. It is easily checked in Boyer–Lindquist coordinates that the ergoregion is described by the inequality $\Delta_r \leq a^2\Delta_\theta \sin^2 \theta$.

Let $\Lambda_\pm \subseteq T^*X$ denote the two components of the conormal bundle to $\{r = r_+\}$,

$$\Lambda_\pm = \{r = r_+, \pm\xi_r > 0, \xi_\theta = \xi_{\phi^*} = 0\},$$

and let $L_\pm$ denote the images in $\partial\overline{T}^*X_+$ of $\Lambda_\pm$ under the canonical projection $T^*X_+ \setminus 0 \rightarrow \partial\overline{T}^*X$. These sets are invariant under the Hamilton flow of p , and $L_\pm \subseteq \widehat{\Sigma}_\pm$. The flow is denoted by

$$\varphi_t = \exp(t \langle \xi \rangle^{-1} H_p)$$

From the dynamical point of view, L_+ is a source and L_- a sink for the Hamilton flow.

Lemma 3.1 ([13, Lemma 4.1]). *There exist neighborhoods $U_\pm$ of $L_\pm$ such that if $(x, \xi) \in U_\pm \setminus L_\pm$, then $\varphi_t(x, \xi) \rightarrow L_\pm$ as $\mp t \rightarrow \infty$ and $\varphi_{\pm T}(x, \xi) \notin U_\pm$ for some $T > 0$.*

To analyze the flow more closely near $r = r_+$, observe that $H_p r$ evaluated at a point $(x, \xi; z)$ where $r = r_+$ is given by

$$H_p r = -g^{-1}(\xi \cdot dx - z dt^*, dr).$$

If $r = r_+$, then

$$\varrho^2 H_p r = (1 - \alpha)(a\xi_{\phi^*} - (r_+^2 + a^2)z).$$

This cannot vanish for $(x, \xi) \in \Sigma \cap T^*X_\delta$ and $z \in \mathbb{R} \setminus 0$, since the condition $p = 0$ implies $\xi_{\phi^*} = a \sin^2 \theta z$. Hence

$$a\xi_{\phi^*} - (r_+^2 + a^2)z = -(r_+^2 + a^2 \cos^2 \theta)z$$

which contradicts $H_p r = 0$. Furthermore, dr is null at $r = r_+$ and

$$g^{-1}(dr, dt^*) = -\varrho^{-2}(1 - \alpha)(r^2 + a^2) < 0$$

at $r = r_+$, so dr lies in the opposite light cone as dt^* . Since Σ_+ is the backwards light cone and Σ_- is the forward light cone, it follows that

$$\pm H_p r < 0 \text{ on } \Sigma_\pm \cap T^*X_\delta \cap \{r = r_+\} \quad (3.1)$$

for $z \in \mathbb{R} \setminus 0$.

Lemma 3.2. *There exists $\delta > 0$ such that φ_t satisfies the following conditions.*

- (1) *If $(x, \xi) \in \widehat{\Sigma}_\pm \setminus L_\pm$, then $\varphi_t(x, \xi) \rightarrow L_\pm$ as $\mp t \rightarrow \infty$, and $\varphi_T(x, \xi) \in \{r \leq r_+ - \delta\}$ for some $\pm T > 0$.*

(2) Suppose $z \in \mathbb{R} \setminus 0$. If $(x, \xi) \in (\Sigma_{\pm} \setminus L_{\pm}) \cap \{|r - r_+| \leq \delta\}$, then either $\varphi_t(x, \xi) \rightarrow L_{\pm}$ as $\mp t \rightarrow \infty$ and

$$\varphi_T(x, \xi) \in \{|r - r_+| \geq \delta\}$$

for some $\pm T > 0$, or

$$\varphi_{\pm T_1}(x, \xi) \in \{r \leq r_+ - \delta\}, \quad \varphi_{\mp T_2}(x, \xi) \in \{r \geq r_+ + \delta\}$$

for some $T_1, T_2 > 0$.

Proof. (1) The first part is proved in Lemma [13, Lemma 4.2], which in turn follows from the same calculations as [30, Section 6.3].

(2) The sets $U_{\pm} \supseteq L_{\pm}$ from Lemma 3.1 have the property that

$$K = (\Sigma \setminus (U_+ \cup U_-)) \cap \{|r - r_+| \leq \delta\}$$

is a compact subset of $\overline{T}^*X_{\delta}$ for $\delta > 0$ sufficiently small. If a flow line enters $U_{\pm}$, then it must tend to $L_{\pm}$ in one direction and leave $U_{\pm}$ in the other according to Lemma 3.1. As it leaves $U_{\pm}$, either $|r - r_+| \geq \delta$ or otherwise it enters the compact set K , at which point $H_p r$ has a definite sign according to (3.1). On the other hand, if a flow line never enters $U_{\pm}$, then it must escape K according to the sign of $H_p r$. $\square$

3.2. Localizing the problem. As remarked in the introduction to this section, an approximate inverse for $P(\lambda)$ is constructed from local inverses, both near the event horizon and near infinity. Begin by decomposing

$$X_{\delta} = \{r_+ - \delta < r < R\} \cup \{r > R/2\},$$

where R will be fixed later. Dealing with the boundaries located at $r = R$ and $r = R/2$ is not convenient, and for this reason these cylinders are embedded in larger manifolds without these boundary components.

Let X_+ denote the larger cylinder $\{r_+ - \delta < r < 3R\}$ capped off with a 3-disk at $\{r = 3R\}$. The operator $P(\lambda)$, defined near $\{r_+ - \delta < r < R\}$, must be extended to X_+ in a suitable way. To do this, define the auxiliary manifold $\mathcal{M}_+ = \mathbb{R}_{t^*} \times X_+$. The goal is to extend the original metric g to $\mathcal{M}_+$, and then consider the stationary Klein–Gordon operator with respect to the extended metric.

The extended metric will be chosen so that $\mathcal{M}_+$ is foliated by spacelike surfaces $\{t^* = \text{constant}\}$. Such a metric is uniquely determined by its ADM decomposition: begin with a smooth function $A > 0$, an arbitrary vector field W , and a Riemannian metric h , all defined initially on X_+ . These objects are extended to $\mathcal{M}_+$ by requiring them to be independent of t^* .

The data (A, W, h) determines a stationary Lorentzian metric g_+ on $\mathcal{M}_+$ by defining

$$\begin{aligned} g_+(T, T) &= A - h(W, W), & g_+(T, V) &= -h(W, V), \\ g_+(V, V) &= -h(V, V), \end{aligned} \tag{3.2}$$

where V is a vector tangent to $\{t^* = \text{constant}\}$.

In standard terminology, A is the lapse function, and W is the shift vector associated to g_+ . If N_t denotes the unit normal to $\{t^* = \text{constant}\}$, then A and W can be uniquely recovered from g_+ via the formulas

$$A = g_+(N_t, T)^2 = g_+^{-1}(dt^*, dt^*)^{-1}, \quad W = \partial_{t^*} - \sqrt{A} N_t.$$

Similarly, h is recovered as the induced metric on $\{t^* = \text{constant}\}$. The idea is to extend g to $\mathcal{M}_+$ by extending the data (A, W, h) originally defined near $\mathbb{R}_{t^*} \times \{r_+ - \delta < r < R\}$.

Lemma 3.3. *Let $R > 0$ be such that T is timelike for g near $\{r > R/10\}$. There exists a stationary Lorentzian metric g_+ on $\mathcal{M}_+$ such that*

- (1) $g_+ = g$ near $\mathbb{R}_{t^*} \times \{r_+ - \delta < r < R\}$,
- (2) dt^* is everywhere timelike for g_+ ,
- (3) T is timelike for g_+ near $\mathbb{R}_{t^*} \times \{r > R/10\}$.

Proof. Let (A, W, h) denote the ADM data for g originally defined near $\mathbb{R}_{t^*} \times \{r_+ - \delta < r < R\}$. Fix any function $\hat{A} > 0$ and Riemannian metric $\hat{h}$ on X_+ such that

$$\hat{A} = A, \quad \hat{h} = h$$

near $\{r_+ - \delta < r < 2R\}$. Choose a cutoff $\chi = \chi(r)$ with values $0 \leq \chi \leq 1$, such that $\text{supp } \chi \subseteq \{r_+ - \delta \leq r < 2R\}$ and $\chi = 1$ near $\{r_+ - \delta \leq r < R\}$. Then, define $\widehat{W} = \chi W$.

The metric g_+ determined by $(\hat{A}, \widehat{W}, \hat{h})$ satisfies the requirements of the lemma; the only part which is not immediate is the last part. If $\chi > 0$, then $A = \hat{A}$ and $h = \hat{h}$, so

$$g_+(T, T) = A - \chi^2 h(W, W) \geq A - h(W, W) > 0.$$

On the other hand, if $\chi = 0$, then $\widehat{W} = 0$ and so $g_+(T, T) = \hat{A} > 0$. □

Given Lemma 3.3, define the stationary operator

$$P_+(\lambda) = e^{i\lambda t^*} (\square_{g_+} + \nu^2 - 9/4) e^{-i\lambda t^*}$$

on X_+ , and similarly for the semiclassical version $P_{h,+}(z)$.

The same argument can also be applied to the infinite part $\{r > R/2\}$: in that case X_∞ is obtained by capping $\{r \geq R/4\}$ at $r = R/4$. Then g is extended to a stationary metric g_∞ on $\mathcal{M}_\infty = \mathbb{R}_{t^*} \times X_\infty$, and the analogue Lemma 3.3 holds in such a way

that T is everywhere timelike for the extended metric. Thus it is possible to define an extended operator $P_\infty(\lambda)$ and $P_{h,\infty}(z)$.

3.3. An approximate inverse near the event horizon. The manifold X_+ can be bordified by adding the artificial boundary

$$H := \{r = r_+ - \delta\}$$

within X_+ . The space of distributions $\overline{H}^1(X_+)$ which are extendible at H is therefore well defined. If $L^2(X_+)$ is defined with respect to any positive density μ on the compact manifold with boundary $X_+ \cup H$, then $\overline{H}^1(X_+)$ agrees with the space of distributions $u \in L^2(X_+)$ such that $du \in L^2(X_+)$ (with respect to any smooth norm on covectors). Then define

$$\mathcal{X}_+ = \{u \in \overline{H}^1(X_+) : P_+(0)u \in L^2(X_+)\},$$

equipped with the norm $\|u\|_{L^2(X_+)} + \|P_+(0)u\|_{L^2(X_+)}$.

Proposition 3.4. *The operator $P_+(\lambda) : \mathcal{X}_+ \rightarrow L^2(X_+)$ is Fredholm of index zero in the half-plane $\{\operatorname{Im} \lambda > -\frac{1}{2}\kappa\}$. Furthermore,*

$$R_+(\lambda) := P_+(\lambda)^{-1} : L^2(X_+) \rightarrow \mathcal{X}_+$$

is a meromorphic family of operators in $\{\operatorname{Im} \lambda > -\frac{1}{2}\kappa\}$ which is holomorphic in any angular sector of the upper half-plane, provided $|\lambda|$ is sufficiently large.

Proof. Let p_+ denote the semiclassical principal symbol of $P_{h,+}(z)$. Note that $p = p_+$ near the characteristic set of p_+ at fiber infinity (whose projection to X_+ is contained in a neighborhood of $\{r < R/10\}$). Therefore the hypotheses at fiber infinity of [30, Section 2.6] also hold for p_+ , since they were verified for p in [13, Section 4]. That suffices to show that $P_+(0)$ is Fredholm on $\mathcal{X}_+$.

The invertibility statement follows from the ellipticity of $P_{h,+}(z)$ on compact subsets of phase space for $\operatorname{Im} z > 0$, which in turn is a corollary of the fact that dt^* is timelike for g_+ — see [30, Sections 3.2, 7]. $\square$

A more refined invertibility statement for $R_+(\lambda)$ will be proved at the end of the section. Before doing this, one must also consider a nontrapping model where $P_{h,+}(z)$ is modified by a complex absorbing operator.

The relevant class of pseudodifferential operators $\Psi_{h,c}^{\operatorname{comp}}(X_+)$ are compactly supported and compactly microlocalized. The compact support means that the Schwartz kernel of any operator in this class is a compact subset of $X_+ \times X_+$, while compactly microlocalized means that the semiclassical wavefront set of any such operator is a compact subset of T^*X_+ . For more about semiclassical microlocal analysis on a non-compact manifold, see [11, Appendix E].

For the purpose of this paper it suffices to observe that any compactly supported function $a \in C_c^\infty(T^*X_+)$ can be quantized to obtain an operator $A = \text{Op}_h(a) \in \Psi_{h,c}^{\text{comp}}(X_+)$ such that $\text{WF}_h(A) \subseteq \text{supp } a$.

In the next lemma, $\Sigma_\pm$ will denote the two components of the characteristic set Σ of p_+ as in Section 3.1 (replacing p with p_+). This separation is possible since dt^* is timelike for g_+ . Furthermore φ_t will denote the Hamilton flow of p_+ .

Lemma 3.5. *Fix $[a, b] \subseteq (0, \infty)$. Then there exists $Q_\pm \in \Psi_{h,c}^{\text{comp}}(X_+)$ such that if $q_\pm = \sigma_h(Q_\pm)$, then for each $z \in [a, b]$,*

- (1) $\text{WF}_h(Q_\pm) \subseteq \Sigma_\pm \cap T^*X_+$ and $\pm q_\pm \leq 0$,
- (2) *If $(x, \xi) \in \Sigma_\pm \setminus L_\pm$, then either $\varphi_t \rightarrow L_\pm$ as $\mp t \rightarrow \infty$, or $\varphi_T(x, \xi) \in \{q_\pm \neq 0\}$ for some $\mp T > 0$.*

Proof. According to Lemma 3.2, there exist compact sets

$$K_\pm \subseteq \Sigma_\pm \cap \{r \geq r_+ + \delta\} \cap T^*X_+$$

such that if $(x, \xi) \in \Sigma_\pm \setminus K_\pm$ then either $\varphi_t(x, \xi) \rightarrow L_\pm$ as $\mp t \rightarrow \infty$, or $\varphi_T(x, \xi) \in K_\pm$ for some $\mp T > 0$. It then suffices to quantize functions $\pm q_\pm \leq 0$ which satisfy $\pm q_\pm < 0$ near $K_\pm$ and have compact support within $\Sigma_\pm \cap T^*X_+$. This may be done uniformly for z in any compact subset of $\mathbb{R} \setminus 0$. $\square$

Fix an interval $[a, b] \subseteq (0, \infty)$ and define $Q = Q_+ + Q_-$, where $Q_\pm$ are given by Lemma 3.5. Then the modified operator $P_{h,+}(z) - iQ$ satisfies the nontrapping condition of [30, Definition 2.12], and hence [30, Theorem 2.14] is valid:

Proposition 3.6. *Fix $[a, b] \subseteq (0, \infty)$ and $[C_-, C_+] \subseteq (-\kappa/2, \infty)$. Then there exists $C > 0$ and $h_0 > 0$ such that*

$$\mathcal{R}_{h,+}(z) := (P_{h,+}(z) - iQ)^{-1} : L^2(X_+) \rightarrow \mathcal{X}_+$$

exists and satisfies the non-trapping bounds

$$\|\mathcal{R}_{h,+}(z)u\|_{L^2(X_+) \rightarrow \overline{H}_h^1(X_+)} \leq Ch^{-1}$$

for $z \in [a, b] + ih[C_-, C_+]$ and $h \in (0, h_0)$.

Remark. *The semiclassical Sobolev space $\overline{H}_h^1(X_+)$ appearing in Proposition 3.6 is $\overline{H}^1(X_+)$ as a set, but equipped with the h -dependent norm*

$$\|u\|_{\overline{H}_h^1(X_+)} = \|u\|_{L^2(X_+)} + h\|du\|_{L^2(X_+)}.$$

The next task is to prove that $R_{h,+}(z)$ exists for $z \in [a, b] + i[Mh, \infty)$ and $M > 0$ sufficiently large, with a corresponding estimate. This does not follow from Proposition 3.4 since there one must take $\text{Im } z > 0$ independent of h . The simplest way to prove this result is by energy estimates.

If h is the induced metric on X_+ , then $|\det g_+| = A \det h$, where $A = g_+^{-1}(dt^*, dt^*)^{-1}$ is the lapse function associated with g_+ . Then differentiating the usual divergence theorem with respect to t^* yields the following: let N_t be the unit normal to X_+ and N_r be the unit normal to $\mathbb{R}_{t^*} \times H$; these are oriented so that N_t points in the direction of increasing t^* , while N_r points in the direction of decreasing r . Then for any vector field V on $\mathcal{M}_+$,

$$\partial_{t^*} \int_{X_+} g_+(V, N_t) dS_t + \int_H g_+(V, N_r) \sqrt{A} dS_H = \int_{X_+} (\operatorname{div}_{g_+} V) \sqrt{A} dS_t. \quad (3.3)$$

Here dS_t is the induced measure on X_+ , and dS_H is the induced measure on H . Note that both N_t and N_r are timelike, and they lie in the same lightcone on their common domain of definition.

The covariant stress-energy tensor $\mathbb{T} = \mathbb{T}[v]$ associated to the Klein–Gordon equation is

$$\mathbb{T}(Y, Z) = [\operatorname{Re}(Yv \cdot Z\bar{v}) - \frac{1}{2}g_+(Y, Z)g_+^{-1}(dv, d\bar{v})] + \frac{1}{2}g_+(Y, Z)(\nu^2 - 9/4)|v|^2.$$

Here v is a sufficiently smooth function on $\mathcal{M}_+$ and Y, Z are real vector fields on $\mathcal{M}_+$. It is well known that the sum in square brackets is positive definite in dv provided Y, Z are both timelike in the same lightcone [22, Lemma 24.1.2].

Given a real vector field Y , let $\mathbb{J}^Y = \mathbb{J}^Y[v]$ be the unique vector field such that $g_+(\mathbb{J}^Y, Z) = \mathbb{T}(Y, Z)$. Then

$$\operatorname{div}_{g_+} \mathbb{J}^Y = \operatorname{Re}((\square_{g_+} + \nu^2 - 9/4)v \cdot Y\bar{v}) + \mathbb{K}^Y(dv, d\bar{v}), \quad (3.4)$$

for a certain tensor $\mathbb{K}^Y$ (the deformation tensor, whose precise properties are not used in this paper). Apply (3.3) to the vector field $\mathbb{J}^Y$. This yields the identity

$$\begin{aligned} \partial_{t^*} \int_{X_+} \mathbb{T}(Y, N_t) dS_t + \int_H \mathbb{T}(Y, N_r) \sqrt{A} dS_H = \\ \int_{X_+} (\operatorname{Re}((\square_{g_+} + \nu^2 - 9/4)v \cdot Y\bar{v}) + \mathbb{K}^Y(dv, d\bar{v})) \sqrt{A} dS_t. \end{aligned} \quad (3.5)$$

Now suppose that Y, Z are stationary in the sense that $[Y, T] = [Z, T] = 0$. Let d_λ denote the covector

$$d_\lambda u := \lambda u dt^* + du,$$

where du is the differential of u on X_+ , and define $Y(\lambda)u := e^{i\lambda t} Y e^{-i\lambda t} u$. Then the stress-energy tensor associated to $e^{-i\lambda t^*} u$ satisfies

$$\begin{aligned} e^{-2\operatorname{Im} \lambda t^*} \mathbb{T}[e^{-i\lambda t^*} u](Y, Z) = \left[\operatorname{Re} \left(Y(\lambda)u \cdot \overline{Z(\lambda)u} \right) - \frac{1}{2}g_+(Y, Z)g_+^{-1}(d_\lambda u, \overline{d_\lambda u}) \right] \\ + \frac{1}{2}(\nu^2 - 9/4)g_+(Y, Z)|u|^2, \end{aligned}$$

and the term inside the square brackets is positive definite in $d_\lambda u$ if Y, Z are timelike in the same light cone.

On the other hand, if $v = e^{-i\lambda t^*} u$ then the integrand on the right hand side of (3.5) can be written as

$$e^{2(\operatorname{Im} \lambda)t^*} \left(\operatorname{Re}(P_+(\lambda)u \cdot \overline{Y(\lambda)u}) + \mathbb{K}^Y(d_\lambda u, \overline{d_\lambda u}) \right).$$

Lemma 3.7. *There exists $C_0, C_1 > 0$ and $C > 0$ such that*

$$|\lambda| \|u\|_{L^2(X_+)} + \|du\|_{L^2(X_+)} \leq \frac{C}{\operatorname{Im} \lambda} \|P_+(\lambda)u\|_{L^2(X_+)}$$

for each $\lambda \in (C_0, \infty) + i(C_1, \infty)$ and $u \in \mathcal{X}_+$.

Proof. Apply (3.5) with the multiplier $Y = N_t$ and $v = e^{-i\lambda t^*} u$. The resulting identity is independent of t^* after multiplying by $e^{-2(\operatorname{Im} \lambda)t^*}$.

Write $f = P_+(\lambda)u$, and consider each term separately. Since N_t is timelike, the zeroth order term in the first integrand on the left can be absorbed by the part which is positive definite in $d_\lambda u$ by taking $|\lambda|$ sufficiently large. The same observation holds for the integral over H , which is positive for $|\lambda|$ sufficiently large. Therefore,

$$\begin{aligned} 2 \operatorname{Im} \lambda \left(|\lambda|^2 \|u\|_{L^2(X_+)}^2 + \|du\|_{L^2(X_+)}^2 \right) \\ \leq \int_{X_+} \left(\operatorname{Re}(f \cdot \overline{N_t(\lambda)u}) + \mathbb{K}^Y(d_\lambda u, \overline{d_\lambda u}) \right) \sqrt{A} dS_t. \end{aligned}$$

Note that $\sqrt{A}$ is bounded by compactness of $X_+ \cup H$, and hence the quadratic form in $d_\lambda u$ coming from the deformation tensor can be absorbed into the left hand side for $\operatorname{Im} \lambda > 0$ sufficiently large. Similarly, Cauchy–Schwarz gives

$$\sqrt{A} |f \cdot \overline{N_t(\lambda)u}| < \frac{|Af|^2}{2\varepsilon \operatorname{Im} \lambda} + \frac{\varepsilon \operatorname{Im} \lambda |N_t(\lambda)u|^2}{2A} \quad (3.6)$$

for $\operatorname{Im} \lambda > 0$, and the second term can be absorbed into the left hand side for $\varepsilon > 0$ sufficiently small. This finally gives

$$2 \operatorname{Im} \lambda \left(|\lambda|^2 \|u\|_{L^2(X_+)}^2 + \|du\|_{L^2(X_+)}^2 \right) \leq \frac{C}{\operatorname{Im} \lambda} \|f\|_{L^2(X_+)}^2$$

as desired. The previous calculations can certainly be justified for $u \in C^\infty(X_+ \cup H)$, which is enough since $C^\infty(X_+ \cup H)$ is dense in $\mathcal{X}_+$ — see [11, Lemma E.4.2] or [30, Section 2.6] for example. $\square$

Remark. By compactness of $X_+ \cup H$, it does not matter how the powers of A are distributed in (3.6). However, this choice is important when proving the analogous result for $P_\infty(\lambda)$ in the next section, noting that the corresponding A in that case satisfies $A \sim r^2$.

Lemma 3.7 implies that $P_+(\lambda)$ is injective for $\lambda \in (C_0, \infty) + i(C_1, \infty)$. Thus $P_+(\lambda)$ is invertible for such λ since it is of index zero according Proposition 3.4. Furthermore,

after applying the semiclassical rescaling, there exists $C > 0$ and $M > 0$ such that for any $[a, b] \subseteq (0, \infty)$,

$$\|R_{h,+}(z)\|_{L^2(X_+) \rightarrow \overline{H}^1_h(X_+)} \leq \frac{C}{Mh} \quad (3.7)$$

for $z \in [a, b] + i[Mh, \infty)$ and h sufficiently small.

3.4. An approximate inverse near infinity. The next step is to prove an analogue of Lemma 3.7 for the operator $P_\infty(\lambda)$ defined in Section 3.2. This will now involve boundary contributions from the conformally timelike boundary $\mathcal{J} = \{r^{-1} = 0\}$, here viewed as a subset of $\mathcal{M}_\infty$. In terms of the boundary defining function $s = r^{-1}$, the metric g_∞ on $\mathcal{M}_\infty$ has an expansion near $\mathcal{J}$ of the form

$$g_\infty = \frac{-ds^2 + \gamma + \mathcal{O}(s^2)}{s^2},$$

where γ is a Lorentzian metric on $\mathcal{J}$. In particular, $\bar{g}_\infty := s^2 g_\infty$ is smooth up to $\mathcal{J}$ and the metric induced on the boundary by $\bar{g}_\infty$ is γ . The function s is extended as a positive function to all of $\mathcal{M}_\infty$ such that $Ts = 0$, and r is extended as well via the formula $r = s^{-1}$.

If $\varepsilon > 0$ is sufficiently small, then $\{s < \varepsilon\}$ defines a neighborhood of $\mathcal{J}$ in $\mathcal{M}_\infty$. Let N_t and N_s be unit normals to X_∞ and $X_\infty \cap \{s = \varepsilon\}$ with respect to g_∞ such that N_t points in the direction of increasing t^* , and N_s points outwards. The analogue of (3.3) becomes

$$\begin{aligned} \partial_{t^*} \int_{X_\infty \cap \{s \geq \varepsilon\}} g_\infty(V, N_t) dS_t - \int_{X_\infty \cap \{s = \varepsilon\}} g_\infty(V, N_s) \sqrt{A} d\mathcal{K} \\ = \int_{X_\infty \cap \{s \geq \varepsilon\}} (\operatorname{div}_{g_\infty} V) \sqrt{A} dS_t \end{aligned} \quad (3.8)$$

Here dS_t and $d\mathcal{K}$ are the measures induced on X_∞ and $X_\infty \cap \{s = \varepsilon\}$ by g_∞ , and A is the lapse function for g_∞ .

The data associated with the foliation by surfaces of constant t^* have conformal analogues: if h is the induced metric on X_+ , then

$$A = s^{-2} \bar{A}, \quad h = s^{-2} \bar{h},$$

where $\bar{A}, \bar{h}$ are smooth up to $\mathcal{J}$. These induce conformally related measures on X_∞ and $X_\infty \cap \{s = \varepsilon\}$ by

$$dS_t = s^{-3} d\bar{S}_t, \quad d\mathcal{K} = s^{-2} d\bar{\mathcal{K}}.$$

In addition $N_t = s \bar{N}_t$ and $N_s = s \bar{N}_s$ where $\bar{N}_t, \bar{N}_s$ are the corresponding unit normals for $\bar{g}_\infty$, smooth up to $\mathcal{J}$. The goal is to eventually let $\varepsilon \rightarrow 0$ in (3.8).

Written in terms of s , the traces $\gamma_\pm$ given by (2.5) have the form

$$\gamma_- v = s^{\nu-3/2} v|_{\mathcal{J}}, \quad \gamma_+ v = -s^{1-2\nu} \partial_s (s^{\nu-3/2} v)|_{\mathcal{J}}.$$

Although $\gamma_{\pm}$ can be given weak formulations, for the energy estimates it is more useful to work with a space of smooth functions on which $\gamma_{\pm}$ are well defined. Given $\nu \in (0, 1)$, let $\mathcal{F}_{\nu}(\mathcal{M}_{\infty})$ denote the space of all $v \in C^{\infty}(\mathcal{M}_{\infty})$ admitting a conormal expansion

$$v(s, y) = s^{3/2+\nu}v_{+}(s^2, y) + s^{3/2-\nu}v_{-}(s^2, y) \quad (3.9)$$

near $\mathcal{I}$, where $(s, y) \in [0, \varepsilon) \times \mathcal{I}$, and $v_{\pm}$ are smooth up to $\mathcal{I}$. If $v \in \mathcal{F}_{\nu}(\mathcal{M}_{\infty})$ is given by (3.9), then

$$\gamma_{-}v(\cdot) = v_{-}(0, \cdot), \quad \gamma_{+}v(\cdot) = (-2\nu)v_{+}(0, \cdot).$$

If $\nu \geq 1$, then one defines $\mathcal{F}_{\nu}(\mathcal{M}_{\infty})$ as all smooth functions on $\mathcal{M}_{\infty}$ vanishing in a neighborhood of $\mathcal{I}$. The space $\mathcal{F}_{\nu}(X_{\infty})$ is defined in the same way, simply replacing $\mathcal{M}_{\infty}$ with X_{∞} .

For general boundary conditions (of the type considered in Section 2.2), the boundary contribution on $X_{\infty} \cap \{s = \varepsilon\}$ arising from the usual stress-energy tensor will diverge as $\varepsilon \rightarrow 0$. This can be remedied by introducing a “twisted” stress-energy tensor as in [20, 31, 32] — the reader is referred to these works for a more complete point of view.

Fix a real, non-vanishing function f . Given a vector field Y on $\mathcal{M}_{\infty}$, define the operator $\tilde{Y}$ by

$$\tilde{Y}v = fY(f^{-1}v),$$

as well as the covector $\tilde{d}v = f d(f^{-1}v)$. The twisted stress-energy tensor $\tilde{\mathbb{T}} = \tilde{\mathbb{T}}[v]$ is defined by

$$\begin{aligned} \tilde{\mathbb{T}}(Y, Z) &= \left[\operatorname{Re} \left(\tilde{Y}v \cdot \tilde{Z}\bar{v} \right) - \frac{1}{2}g_{\infty}(Y, Z)g_{\infty}^{-1}(\tilde{d}v, \tilde{d}\bar{v}) \right] \\ &\quad + \frac{1}{2}g_{\infty}(Y, Z)(F + \nu^2 - 9/4)|v|^2, \end{aligned} \quad (3.10)$$

where $F = f^{-1}\square_{g_{\infty}}(f)$ is a scalar potential term. The term in square brackets is positive definite in $\tilde{d}v$ provided Y, Z are timelike in the same lightcone. The twisting function f is chosen so that

$$F + \nu^2 - 9/4 = \mathcal{O}(s^2).$$

If Y, Z are smooth up to $\mathcal{I}$, this guarantees that $g_{\infty}(Y, Z)(F + \nu^2 - 9/4)$ is also smooth up to $\mathcal{I}$. The simplest choice of f with this property is $f = s^{3/2-\nu}$.

Remark. Since $(F + \nu^2 - 9/4)|v|^2$ is of zeroth order in v , the precise sign properties of F will not be important in the high frequency regime (just as in Lemma 3.7). More refined choices of f leading to positive F are discussed at length in [20].

Next, let $\tilde{\mathbb{J}}^Y = \tilde{\mathbb{J}}^Y[v]$ denote the associated energy current, namely $g_{\infty}(\tilde{\mathbb{J}}^Y, Z) = \tilde{\mathbb{T}}(Y, Z)$.

Lemma 3.8. *Suppose that Y Killing for g_∞ and $Yf = 0$. Then*

$$\operatorname{div}_{g_\infty} \tilde{\mathbb{J}}^Y = \operatorname{Re} \left((\square_{g_\infty} + \nu^2 - 9/4)v \cdot Y\bar{v} \right)$$

Proof. Since Y is Killing, the condition $Yf = 0$ also implies $YF = 0$, and then the result follows from a direct calculation [32, Lemma 2.5]. $\square$

If $f = s^{3/2-\nu}$, then $Tf = 0$ and hence Lemma 3.8 is valid. Now apply (3.8) to the vector field $\tilde{\mathbb{J}}^T$, where $v \in \mathcal{F}_\nu(\mathcal{M}_\infty)$. Consider the integral over $X_\infty \cap \{s \geq \varepsilon\}$, which can be written as

$$\int_{X_\infty \cap \{s \geq \varepsilon\}} \tilde{\mathbb{T}}(T, \bar{N}_t) s^{-2} d\bar{S}_t.$$

Checking the various powers of s , this integral has a limit as $\varepsilon \rightarrow 0$ for $v \in \mathcal{F}_\nu(\mathcal{M}_\infty)$. This also motivates the following spaces: let $\mathcal{L}^2(X_\infty)$ denote the space of distributions for which

$$\|u\|_{\mathcal{L}^2(X_\infty)} = \int_{X_\infty} |u|^2 s dS_t < \infty,$$

and let $\mathcal{H}^1(X_\infty)$ denote the space of distributions for which

$$\|u\|_{\mathcal{H}^1(X_\infty)} = \int_{X_\infty} \left(|u|^2 + s^{-2} g_\infty^{-1}(\tilde{d}u, \tilde{d}\bar{u}) \right) s dS_t < \infty.$$

Compare these spaces with those defined in Section 2.2.

Next, consider the integral in (3.8) over $X_\infty \cap \{s = \varepsilon\}$. This is only relevant in the case $\nu \in (0, 1)$ since if $\nu \geq 1$ then the integral automatically vanishes for $v \in \mathcal{F}_\nu(\mathcal{M}_\infty)$ and $\varepsilon > 0$ sufficiently small. Since T and N_s are orthogonal and $Tf = 0$, this reduces to

$$\int_{X_\infty \cap \{s = \varepsilon\}} \operatorname{Re} \left(Tv \cdot \tilde{N}_s \bar{v} \right) s^{-3} \bar{A}^{1/2} d\bar{K}. \quad (3.11)$$

Furthermore, $N_s = -s\partial_s + \mathcal{O}(s^3)$. Write

$$s^{-3} Tv \cdot \tilde{N}_s \bar{v} = (s^{\nu-3/2} Tv) \left(s^{-3/2-\nu} \tilde{N}_s \bar{v} \right),$$

and notice that this tends to $\gamma_-(Tv) \cdot \gamma_+ \bar{v}$ as $\varepsilon \rightarrow 0$ for $v \in \mathcal{F}_\nu(X_\infty)$.

Taking $\varepsilon \rightarrow 0$ in (3.8), one therefore has the identity

$$\begin{aligned} \partial_{t^*} \int_{X_\infty} \tilde{\mathbb{T}}(T, N_t) dS_t - \int_{X_\infty \cap \mathcal{I}} \operatorname{Re} (\gamma_-(Tv) \cdot \gamma_+ \bar{v}) \bar{A}^{1/2} d\bar{K} \\ = \int_{X_\infty} \operatorname{Re} \left((\square_{g_\infty} + \nu^2 - 9/4)v \cdot T\bar{v} \right) \sqrt{A} dS_t. \end{aligned} \quad (3.12)$$

Using (3.12), it is now straightforward to prove the analogue of Lemma 3.7. In the following, either $B = \gamma_-$ or $B = \gamma_+ + \beta\gamma_-$, where $\beta \in C^\infty(\mathcal{I}; \mathbb{R})$ satisfies $T\beta = 0$; therefore β may be considered as a function on $X_\infty \cap \mathcal{I}$.

Lemma 3.9. *There exists $C_0 > 0$ and $C > 0$ such that*

$$|\lambda| \|u\|_{\mathcal{L}^2(X_\infty)} + \|u\|_{\mathcal{H}^1(X_\infty)} \leq \frac{C}{\operatorname{Im} \lambda} \|P_\infty(\lambda)u\|_{r^{-2}\mathcal{L}^2(X_\infty)}$$

for each $\lambda \in (C_0, \infty) + i(0, \infty)$ and $u \in \mathcal{F}_\nu(X_\infty)$ such that $Bu = 0$ if $\nu \in (0, 1)$.

Proof. The proof is close to that of Lemma 3.7. Apply (3.12) to a function $v = e^{-i\lambda t^*}u$, where $u \in \mathcal{F}_\nu(X_\infty)$, and multiply the resulting identity by $e^{-2\operatorname{Im} \lambda t^*}$. Since T and $\bar{N}_t$ are timelike in the same lightcone, the first integral in (3.12) controls

$$2\operatorname{Im} \lambda \left(|\lambda|^2 \|u\|_{\mathcal{L}^2(X_\infty)}^2 + \|u\|_{\mathcal{H}^1(X_\infty)}^2 \right)$$

for $\operatorname{Im} \lambda > 0$ and $|\lambda|$ sufficiently large. Write $f = (\square_g + \nu^2 - 9/4)u$, and write the integral on the right hand side of (3.12) in terms of the conformal measure. Then the integrand is bounded by

$$s^{-4} \bar{A}^{1/2} \operatorname{Re}(i\bar{\lambda}u \cdot f) \leq \frac{s^{-2}}{2\varepsilon \operatorname{Im} \lambda} \cdot |s^{-2}f|^2 + \frac{\varepsilon s^{-2} \operatorname{Im} \lambda |\lambda|^2}{2} \bar{A} \cdot |u|^2. \quad (3.13)$$

Now integrate (3.13) over X_∞ with respect to $d\bar{S}_t$. The integral of the first term on the right hand side is bounded by $C(\operatorname{Im} \lambda)^{-1} \|f\|_{r^{-2}\mathcal{L}^2(X_\infty)}^2$, while the integral of the second term can be absorbed into the left hand side for ε sufficiently small.

It remains to handle the integral over $X_\infty \cap \mathcal{J}$. If u satisfies Dirichlet boundary condition (which recall is automatic for $\nu \geq 1$), then this term vanishes. Otherwise, if $\nu \in (0, 1)$ and $B = \gamma_+ + \beta\gamma_-$, then the integrand becomes (after accounting for the minus sign in (3.12))

$$2\operatorname{Im} \lambda \int_{X_\infty \cap \mathcal{J}} \beta \cdot |\gamma_- u|^2 \bar{A}^{1/2} d\bar{\mathcal{K}}.$$

If β is nonnegative this term can be dropped. Otherwise, note that for each $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that

$$\int_{X_\infty \cap \mathcal{J}} |\gamma_- u|^2 d\bar{\mathcal{K}} \leq \varepsilon \|u\|_{\mathcal{H}^1(X_\infty)}^2 + C_\varepsilon \|u\|_{\mathcal{L}^2(X_\infty)}^2.$$

This is proved directly in [32, Lemma B.1]; alternatively, it also follows from the boundedness of the trace $\gamma_- : \mathcal{H}^1(X_\infty) \rightarrow H^\nu(X_\infty \cap \mathcal{J})$ and the compactness of the inclusion $H^\nu(X_\infty \cap \mathcal{J}) \hookrightarrow L^2(X_\infty \cap \mathcal{J})$. Hence the boundary term can always be absorbed by the left hand side for any $\operatorname{Im} \lambda > 0$ and $|\lambda|$ sufficiently large. $\square$

Now define

$$\mathcal{X}_\infty = \begin{cases} \{u \in \mathcal{H}^1(X_\infty) : P_\infty(0)u \in r^{-2}\mathcal{L}^2(X_\infty)\} & \text{if } \nu \geq 1, \\ \{u \in \mathcal{H}^1(X_\infty) : P_\infty(0)u \in r^{-2}\mathcal{L}^2(X_\infty) \text{ and } Bu = 0\} & \text{if } \nu \in (0, 1), \end{cases} \quad (3.14)$$

equipped with the graph norm — compare to Section 2.2. Note that $P_\infty(\lambda)$ is elliptic at fiber-infinity $\partial \bar{T}^* X_\infty$, where X_∞ viewed as a noncompact manifold without boundary. Furthermore, $P_\infty(\lambda)$ is elliptic at $X_\infty \cap \mathcal{J}$ as a Bessel operator with respect to the

boundary defining function $s = r^{-1}$. By elliptic regularity [15, Theorem 1 and Lemma 4.13] it is therefore possible to show that all computations in Lemma 3.9 for $u \in \mathcal{F}_\nu(X_\infty)$ are also valid for $u \in \mathcal{X}_\infty$. In particular,

$$|\lambda| \|u\|_{\mathcal{L}^2(X_\infty)} + \|u\|_{\mathcal{H}^1(X_\infty)} \leq \frac{C}{\operatorname{Im} \lambda} \|P_\infty(\lambda)u\|_{r^{-2}\mathcal{L}^2(X_\infty)} \quad (3.15)$$

for each $u \in \mathcal{X}_\infty$ and $\lambda \in (C_0, \infty) + i(0, \infty)$.

In addition, since B is an elliptic boundary condition in the sense of Bessel operators [15, Section 4.4], one has the following:

Lemma 3.10. *The operator $P_\infty(\lambda) : \mathcal{X}_\infty \rightarrow r^{-2}\mathcal{L}^2(X_\infty)$ is Fredholm of index zero, and is invertible outside arbitrarily small angles about the real axis for $|\lambda|$ sufficiently large.*

As a corollary of (3.15), the operator $R_\infty(\lambda) := P_\infty(\lambda)^{-1}$ exists for $\lambda \in (C_0, \infty) + i(0, \infty)$. In terms of the semiclassical rescaling, there exists $C > 0$ such that for each $[a, b] \subseteq (0, \infty)$,

$$\|R_{h,\infty}(z)\|_{r^{-2}\mathcal{L}^2(X_\infty) \rightarrow \mathcal{H}_h^1(X_\infty)} \leq \frac{C}{\operatorname{Im} z} \quad (3.16)$$

for $z \in [a, b] + i(0, \infty)$ and h sufficiently small.

3.5. Construction a global approximate inverse. Combining the results of Sections 3.3, 3.4, it is now possible to construct a global approximate inverse for $P(\lambda)$ on X_δ .

Fix a smooth partition of unity $\chi_1 + \chi_2 = 1$ and functions ψ_1, ψ_2 on X_δ such that

- $\operatorname{supp} \chi_1 \cup \operatorname{supp} \psi_1 \subseteq \{r > R/2\}$ and $\operatorname{supp} \chi_2 \cup \operatorname{supp} \psi_2 \subseteq \{r < R\}$,
- $\psi_1 = 1$ near $\operatorname{supp} \chi_1$ and $\psi_2 = 1$ near $\operatorname{supp} \chi_2$.

Begin by fixing $M > 0$ such that (3.7) holds. Then, choose $[C_-, C_+] \subseteq (-\kappa/2, \infty)$ such that $M \in (C_-, C_+)$ and $|C_+ - M| \geq |C_- - M|$, increasing C_+ if necessary.

Given an interval $[a, b] \subseteq (0, \infty)$, the operators $\mathcal{R}_{h,+}(z), R_{h,+}(z_0), R_{h,\infty}(z_0)$ exist for $z \in [a, b] + ih[C_-, C_+]$ and $z_0 \in [a, b] + ih[M, C_+]$. Define

$$\begin{aligned} E(z, z_0) &= \psi_1 R_{h,\infty}(z_0) \chi_1 \\ &\quad + \psi_2 (\mathcal{R}_{h,+}(z) - i R_{h,+}(z_0) Q \mathcal{R}_{h,+}(z)) \chi_2, \end{aligned}$$

where $z \in [a, b] + ih[C_-, C_+]$ and $z_0 \in [a, b] + ih[M, C_+]$. This is a well defined operator $r^{-2}\mathcal{L}^2(X_\delta) \rightarrow \mathcal{X}$ in view of the cutoffs, and it is holomorphic in z for each z_0 .

Apply $P_h(z)$ to $E(z, z_0)$ on the left: the first term yields

$$\chi_1 + \psi_1 (P_{h,\infty}(z) - P_{h,\infty}(z_0)) R_{h,\infty}(z_0) \chi_1 + [P_h(z), \psi_1] R_{h,\infty}(z_0) \chi_1, \quad (3.17)$$

while the second term yields

$$\begin{aligned} \chi_2 - i\psi_2 (P_{h,+}(z) - P_{h,+}(z_0)) R_{h,+}(z_0) Q \mathcal{R}_{h,+}(z) \chi_2 \\ + [P_h(z), \psi_2] (\mathcal{R}_{h,+}(z) - iR_{h,+}(z_0) Q \mathcal{R}_{h,+}(z)) \chi_2. \end{aligned} \quad (3.18)$$

Adding (3.17), (3.18), one has

$$P_h(z)E(z, z_0) = I + K_1(z, z_0) + K_2(z, z_0) + K_3(z, z_0) + K_4(z, z_0),$$

where K_1, K_2 are the second and third terms in (3.17), and K_3, K_4 are the second and third terms in (3.18). Also let $K = K_1 + K_2 + K_3 + K_4$.

Lemma 3.11. *There exist compactly supported pseudodifferential operators*

$$A(z) \in h\Psi_{h,c}^{-1}(X_+), \quad B(z) \in h\Psi_{h,c}^{-\infty}(X_+)$$

depending smoothly on z such that

$$[P_h(z), \psi_2] \mathcal{R}_{h,+}(z) = A(z) + B(z) \mathcal{R}_{h,+}(z)$$

for $z \in [a, b] + ih[C_-, C_+]$.

Proof. The commutator $[P_{h,+}(z), \psi_2]$ has coefficients supported near $\text{supp } d\psi_2$ where it may be assumed that $P_{h,+}(z)$ is elliptic at fiber infinity $\partial \bar{T}^* X_+$. Choose $\varphi, \varphi' \in C_c^\infty(X_+)$ satisfying $\varphi = 1$ near $\text{supp } d\phi_2$ and $\varphi' = 1$ near $\text{supp } \varphi$. By choosing φ' with sufficiently small support it may be assumed that $P_{h,+}(z)$ is elliptic near $\pi^{-1}(\text{supp } \varphi)$, where $\pi : \partial \bar{T}^* X_+ \rightarrow X_+$ is the canonical projection. By ellipticity, there exist properly supported operators

$$F(z) \in \Psi_h^{-2}(X_+), \quad Y(z) \in h^\infty \Psi_h^{-\infty}(X_+)$$

and $C \in \Psi_{h,c}^{\text{comp}}(X_+)$ such that

$$\varphi = F(z)\varphi' P_{h,+}(z) + Y(z) + C.$$

The operators $F(z), Y(z)$ may be chosen to depend smoothly on z , and C can be chosen uniformly for z in a compact set. Then,

$$\begin{aligned} [P_h(z), \psi_2] \mathcal{R}_{h,+}(z) &= [P_h(z), \psi_2] \varphi \mathcal{R}_{h,+}(z) \\ &= [P_h(z), \psi_2] (F(z)\varphi' + (Y(z) + C) \mathcal{R}_{h,+}(z)). \end{aligned}$$

Since the commutator lies in $h\Psi_h^1(X_+)$, it suffices to define $A(z) = [P_h(z), \psi_2] F(z)\varphi'$ and $B(z) = [P_h(z), \psi_2] (Y(z) + C)$. $\square$

Lemma 3.12. *If $M > 0$ is sufficiently large, then there exists h_0 depending on M such that the following hold for $z \in [a, b] + ih[C_-, C_+]$ and $z \in [a, b] + ih[M, C_+]$.*

(1) $K(z, z_0) : r^{-2} \mathcal{L}^2(X_\delta) \rightarrow r^{-2} \mathcal{L}^2(X_\delta)$ is compact.

(2) $I + K(z_0, z_0)$ is invertible for $h \in (0, h_0)$.

Proof. To prove compactness, consider each term in $K(z, z_0)$ separately. For K_1, K_2 , the operator $P_{h,\infty}(z)$ is an elliptic Bessel operator (in the non-semiclassical sense) [15, Section 4.4]. Furthermore, if $\nu \in (0, 1)$, then the boundary operator B is elliptic in the sense of [15, Section 4.4]. By elliptic regularity [15, Theorem 1],

$$(P_{h,+}(z) - P_{h,+}(z_0)) R_{h,+}(z_0) : r^{-2}\mathcal{L}^2(X_+) \rightarrow r^{-2}\mathcal{H}^1(X_+)$$

is bounded, noting that $P_{h,+}(z) - P_{h,+}(z_0)$ is of first order. The inclusion $r^{-2}\mathcal{H}^1(X_+) \hookrightarrow r^{-2}\mathcal{L}^2(X_+)$ is compact, which shows that

$$K_1(z, z_0) : r^{-2}\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{L}^2(X_\delta)$$

is compact. A similar argument also shows that $K_2(z, z_0)$ is compact.

For $K_3(z, z_0), K_4(z, z_0)$, each of the terms containing Q are compact since any operator in $\Psi_{h,c}^{\text{comp}}(X_+)$ is smoothing. It remains to consider the commutator term $[P_h(z), \psi_2]\mathcal{R}_{h,+}(z)$. For that, compactness follows from Lemma 3.11.

To prove the invertibility statement, notice that for $z = z_0$,

$$K(z_0, z_0) = [P_h(z_0), \psi_1]R_{h,\infty}(z_0)\chi_1 + [P_h(z_0), \psi_2]R_{h,+}(z_0)\chi_2. \quad (3.19)$$

As first order operators the commutators are of order $\mathcal{O}(h)$, where the implicit constants are independent of any $M > 0$ for h sufficiently small depending on M . By choosing $M > 0$ sufficiently large and applying (3.7), (3.16), the operator norm of $K(z_0, z_0)$ is of order $\mathcal{O}(M^{-1})$, hence $I + K(z_0, z_0)$ is invertible by Neumann series for h sufficiently small. $\square$

From now on it will be assumed that $M > 0$ is chosen sufficiently large so that Lemma 3.12 holds. This can always be achieved *before* selecting $C_\pm$ since (3.19) does not involve the operator $\mathcal{R}_{h,+}(z)$.

Since $K(z, z_0)$ is compact and $I + K(z_0, z_0)$ is invertible for an appropriate choice of z_0 with h small, it follows that $(I + K(z, z_0))^{-1} : r^{-2}\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{L}^2(X_\delta)$ is a meromorphic family of operators. If $(I + K(z, z_0))^{-1}$ exists, then $P_h(z) : \mathcal{X} \rightarrow r^{-2}\mathcal{L}^2(X_\delta)$ has a right inverse given by $E(z, z_0)(I + K(z, z_0))^{-1}$. In that case $P_h(z)$ is invertible, since it is of index zero by Theorem 2. Analytic continuation then shows that

$$R_h(z) = E(z, z_0)(I + K(z, z_0))^{-1}.$$

Furthermore, any pole of $R_h(z)$ is also a pole of $(I + K(z, z_0))^{-1}$.

3.6. Singular values. In order to prove (2.3) of Proposition 2.3, one must bound

$$\begin{aligned} \|R_h(z)\|_{r^{-2}\mathcal{L}^2(X_\delta) \rightarrow \mathcal{L}^2(X_\delta)} \\ \leq \|E(z, z_0)\|_{r^{-2}\mathcal{L}^2(X_\delta) \rightarrow \mathcal{L}^2(X_\delta)} \|(I + K(z, z_0))^{-1}\|_{r^{-2}\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{L}^2(X_\delta)}. \end{aligned}$$

Using (3.7), (3.16) and Lemma 3.2, the operator norm of $E(z, z_0)$ is of order $\mathcal{O}(h^{-2})$, which will be harmless compared to the exponentially growing bound on the norm of $(1 + K(z, z_0))^{-1}$.

Lemma 3.13 ([16, Theorem V.5.1]). *Let $\mathcal{H}$ be a Hilbert space. Then*

$$\|(I + A)^{-1}\|_{\mathcal{H} \rightarrow \mathcal{H}} \leq \frac{\det(I + |A|)}{|\det(I + A)|}.$$

for any operator $A : \mathcal{H} \rightarrow \mathcal{H}$ of trace class.

Lemma 3.13 cannot be applied directly to $I + K(z, z_0)$ since $K(z, z_0)$ is not of trace class. Instead, $K(z, z_0)$ lies in a Schatten p -class for some $p > 0$. For a compact operator $A : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ between Hilbert spaces, let $s_j(A) = s_j(A; \mathcal{H}_1, \mathcal{H}_2)$, $j \in \mathbb{N}_{\geq 1}$ denote its singular values, listed in decreasing order with multiplicity.

Lemma 3.14. *There exists $C > 0$ such that the singular values of $K(z, z_0)$ satisfy*

$$s_j(K(z, z_0)) \leq Ch^{-3} j^{-1/3}$$

uniformly for $z \in [a, b] + ih[C_-, C_+]$ and $z_0 \in [a, b] + ih[M, C_+]$.

Proof. By Fan's inequality $s_{i+j-1}(A + B) \leq s_i(A) + s_j(B)$ applied repeatedly,

$$s_j(K(z, z_0)) \leq \sum_{i=1}^4 s_{4j-3}(K_i(z, z_0)).$$

The operator $K_1(z, z_0) : r^{-2}\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{H}_h^1(X_\infty)$ is of order $\mathcal{O}(h^{-1})$, whereas the inclusion $\mathcal{H}_h^1(X_\infty) \hookrightarrow \mathcal{L}^2(X_\infty)$ has singular values bounded by $Ch^{-1}j^{-1/3}$ [15, Appendix B]. By standard properties of singular values,

$$s_j(K_1(z, z_0)) \leq Ch^{-2}j^{-1/3}.$$

The same argument applies to $K_2(z, z_0)$, so accounting for the extra power of h coming from the commutator,

$$s_j(K_1(z, z_0)) \leq Ch^{-1}j^{-1/3}.$$

The terms K_2, K_3 can be handled similarly, using that the inclusion $\overline{H}_h^1(X_+) \hookrightarrow L^2(X_+)$ has singular values bounded by $Ch^{-1}j^{-1/3}$. The only term that requires extra care is $[P_h(z), \psi_2]\mathcal{R}_{h,+}(z)$ in K_4 , but by using Lemma 3.11 this operator is seen to map $L^2(X_+) \rightarrow \overline{H}_h^1(X_+)$ with operator norm of order $\mathcal{O}(h)$. $\square$

It follows from Lemma 3.14 that $K(z, z_0)^4$ is of trace class, and using Fan's inequality $s_{i+j-1}(AB) \leq s_i(A)s_j(B)$, one has the estimate

$$s_j(K(z, z_0)^4) \leq s_{4j-3}(K(z, z_0))^4 \leq Ch^{-12}j^{-4/3}. \quad (3.20)$$

This is uniform for $z \in [a, b] + ih[C_-, C_+]$ and $z_0 \in [a, b] + ih[M, C_+]$. To apply Lemma 3.13, write

$$(I + K(z, z_0))^{-1} = \left(\sum_{j=0}^3 (-1)^j K(z, z_0)^j \right) (I - K(z, z_0)^4)^{-1} \quad (3.21)$$

The norm of $K(z, z_0) : r^{-2}\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{L}^2(X_\delta)$ is of order $\mathcal{O}(h^{-2})$, so the norm of the sum on the right hand side of (3.21) is polynomially bounded in h .

Note from (3.21) that any pole of $(I + K(z, z_0))^{-1}$ is a pole of $(I - K(z, z_0)^4)^{-1}$, hence the poles of $R_h(z)$ are among those of $(I - K(z, z_0)^4)^{-1}$. Now Lemma 3.13 is applied to $K(z, z_0)^4$.

Lemma 3.15. *Let $F(h)$ denote the supremum of $\det(I + |K(z, z_0)^4|)$ for $z \in [a, b] + ih[C_-, C_+]$ and $z_0 \in [a, b] + ih[M, C_+]$. Then*

$$F(h) \leq e^{Ch^{-12}}$$

for some $C > 0$.

Proof. The determinant is bounded by

$$\det(I + |K(z, z_0)^4|) = \prod_{j \geq 1} (1 + s_j(K(z, z_0)^4)) \leq \exp \left(Ch^{-12} \sum_{j \geq 1} j^{-4/3} \right)$$

according to (3.20). □

The next step is to bound $|f(z, z_0)|$ from below, where $f(z, z_0) = \det(I - K(z, z_0)^4)$.

Lemma 3.16. *The function $f(z, z_0)$ has the following properties.*

- (1) $|f(z, z_0)| \leq F(h)$.
- (2) $f(z_0, z_0) \neq 0$, and moreover $|f(z_0, z_0)| \geq e^{-Ch^{-12}}$ for some $C > 0$.

Proof. 1) The estimate $|f(z, z_0)| \leq F(h)$ follows from Weyl convexity inequalities [11, Proposition B.2.4].

2) As in the proof of Lemma 3.12, the norm of $K(z_0, z_0)^4$ is of order $\mathcal{O}(M^{-4})$ for h sufficiently small. By increasing $M > 0$ if necessary, the operator $I + K(z_0, z_0)^4$ is invertible, and

$$(I - K(z_0, z_0)^4)^{-1} = I + K(z_0, z_0)^4(I - K(z_0, z_0)^4)^{-1}. \quad (3.22)$$

Arguing as in Lemma 3.15,

$$\det(I + |K(z_0, z_0)^4(I + K(z_0, z_0)^4)^{-1}|) \leq e^{Ch^{-12}},$$

which gives $|f(z_0, z_0)| \geq e^{-Ch^{-12}}$. □

The proof of Proposition 2.3 can now be finished using the following lemma of Cartan [24, Theorem 11]:

Lemma 3.17. *Suppose that $g(z)$ is holomorphic in a neighborhood of a disk $B(z_0, R)$, such that $g(z_0) \neq 0$. Fix $r \in (0, R)$, and let $\{w_j\}$ denote the zeros of $g(z)$ in $B(z_0, R)$ for $j = 1, \dots, n(z_0, R)$. Given any $\rho > 0$,*

$$\log |g(z)| - \log |g(z_0)| \geq -\frac{2r}{R-r} \log \left(\sup_{|z-z_0| < R} |g(z)| \right) - n(z_0, R) \log \left(\frac{(R+r)}{\rho} \right)$$

for $z \in B(z_0, R) \setminus \bigcup_j B(w_j, \rho)$.

Lemma 3.17 will be applied to the function $z \mapsto f(z, z_0)$ and disks of radius proportional to h . This requires a bound on the number of zeros of $f(z, z_0)$ in disks of the form $B(z_0, Rh)$.

As noted at the beginning of Section 3.5, it may always be assumed that $|C_+ - M| \geq |C_- - M|$. Recall the definitions of $\Omega_\varepsilon(h)$ and $\Omega(h)$ from (2.10). Then given $\varepsilon > 0$ there exists $M' \geq M$ and $R > 0$ such that the union of all disks $B(w, Rh)$ with $w \in [a, b] + ih[M, M']$ covers $\Omega(h)$ and is contained in $\Omega_\varepsilon(h)$. If $\varepsilon > 0$ is sufficiently small, then

$$\Omega_{3\varepsilon}(h) \subseteq [a', b'] + ih[C'_-, C'_+],$$

where $[a', b'] \subseteq (0, \infty)$ and $[C'_-, C'_+] \subseteq (-\kappa/2, \infty)$. Applying Lemmas 3.15, 3.16 to this larger rectangle shows that certainly $|f(z, z_0)| \leq e^{Ch^{-12}}$ for $z_0 \in [a, b] + i[M, M']$ and $z \in B(z_0, (R + 2\varepsilon)h)$.

Lemma 3.18. *Let $z_0 \in [a, b] + ih[M, M']$. Then there exists $C > 0$ such*

$$n(z_0, (R + \varepsilon)h) \leq Ch^{-12},$$

uniformly in z_0 , where $n(z_0, (R + \varepsilon)h)$ is the number of zeros of $f(z, z_0)$ in $B(z_0, (R + \varepsilon)h)$.

Proof. By Jensen's formula, the number of zeros $n(z_0, \rho)$ of $f(z, z_0)$ within $B(z_0, \rho)$ satisfies

$$\begin{aligned} \int_0^{h(R+2\varepsilon)} \frac{n(z_0, \rho)}{\rho} d\rho &= \frac{1}{2\pi} \int_0^{2\pi} \log |f(z_0 + h(R+2\varepsilon)e^{i\theta}, z_0)| d\theta - \log |f(z_0, z_0)| \\ &\leq Ch^{-12}. \end{aligned}$$

Therefore the number of zeros in a disk $n(z_0, (R + \varepsilon)h)$ is estimated by

$$\frac{R+2\varepsilon}{\varepsilon} n(z_0, (R + \varepsilon)h) \leq \int_{h(R+\varepsilon)}^{h(R+2\varepsilon)} \frac{n(z_0, \rho)}{\rho} d\rho \leq Ch^{-12}.$$

□

Remark. For each fixed z_0 , the poles of $R_h(z)$ are among the zeros of $f(z, z_0)$. The region $\Omega(h)$ can be covered by at most $\mathcal{O}(h^{-1})$ disks of radius Rh with centers in $[a, b] + ih[M, M']$. According to Lemma 3.18, $R_h(z)$ has at most $\mathcal{O}(h^{-12})$ poles in each of these disks, so altogether $R_h(z)$ has at most $\mathcal{O}(h^{-13})$ poles in $\Omega(h)$.

Combining Lemma 3.17 with Lemma 3.18 shows that for any function $0 < S(h) = o(h)$ and $z_0 \in [a, b] + ih[M, M']$,

$$|f(z, z_0)| \geq \exp(-Ch^{-12} \log(1/S(h))), \quad z \in B(z_0, Rh) \setminus \bigcup_j B(w_j, S(h)), \quad (3.23)$$

where $\{w_j\}$ are the zeros of $f(z, z_0)$ in $B(z_0, (R + \varepsilon)h)$.

Proof of Proposition 2.3. Combine Lemma 3.15 with (3.23). This shows that for any $z_0 \in [a, b] + ih[M, M']$,

$$\|R_h(z)\|_{r^{-2}\mathcal{L}^2(X_\delta) \rightarrow r^{-2}\mathcal{L}^2(X_\delta)} \leq \exp(Ch^{-12} \log(1/S(h))) \quad (3.24)$$

for

$$z \in B(z_0, Rh) \setminus \bigcup_j B(w_j, S(h)),$$

where $\{w_j\}$ are the zeros of $f(z, z_0)$ in $B(z_0, (R + \varepsilon)h)$. If w_j is not a pole of $R_h(z)$, then apply the maximum principle to the holomorphic operator-valued function $R_h(z)$ on $B(w_j, S(h))$ to see that (3.24) is valid on $B(w_j, S(h))$ as well. Thus (3.24) holds for $z \in B(z_0, Rh) \setminus \bigcup_j B(z_j, S(h))$, where now $\{z_j\}$ denote the poles of $R_h(z)$ in $B(z_0, (R + \varepsilon)h)$.

The disks $B(z_0, Rh), B(z_0, (R + \varepsilon)h)$ for $z_0 \in [a, b] + ih[M, M']$ cover $\Omega(h), \Omega_\varepsilon(h)$ respectively, whence the result follows. $\square$

4. PROOF OF PROPOSITION 2.1

To complete the proof of Theorem 1 it remains to prove Proposition 2.1. The proof is very similar to that of Lemmas 3.7 and 3.9. The twisted stress-energy tensor $\tilde{\mathbb{T}} = \tilde{\mathbb{T}}[v]$ is defined as in (3.12) using the metric g ; the twisting function is again $f = r^{\nu-3/2}$.

The future pointing Killing generator K of the null surface $\{r = r_+\}$ can be normalized by requiring $Kt^* = 1$, in which case

$$K = T + \frac{a}{r_+^2 + a^2} \Phi.$$

Let $d\sigma$ denote the measure induced on $X_0 \cap \{r = r_+\}$. With the above normalization, the analogue of (3.12) on X_0 has the form

$$\begin{aligned} \partial_{t^*} \int_{X_0} \tilde{\mathbb{T}}(Y, N_t) dS_t - \int_{X_0 \cap \mathcal{H}} \operatorname{Re}(\gamma_- (Yv) \cdot \gamma_+ \bar{v}) \bar{A}^{1/2} d\bar{K} \\ = - \int_{X_0 \cap \{r=r_+\}} \tilde{\mathbb{T}}(Y, K) \sqrt{A} d\sigma + \int_{X_0} \operatorname{Re}((\square_g + \nu^2 - 9/4)v \cdot Y\bar{v}) \sqrt{A} dS_t, \end{aligned}$$

where Y is a Killing field such that $Yr = 0$ and $A = g^{-1}(dt^*, dt^*)^{-1}$ is the lapse function.

This identity is applied with the vector field $Y = K$. The contribution from the horizon is the integral of

$$\tilde{\mathbb{T}}(K, K) = |Kv|^2 \geq 0,$$

which may be dropped in view of its nonnegativity.

If $|a| < r_+^2$, then K is everywhere timelike on $\mathcal{M}_0$. In that case Proposition 2.1 would follow from the same proof as in Lemma 3.9; the only difference is that coercivity of the derivative transverse to the horizon degenerates at the horizon. This does not affect the final result since Proposition 2.1 only involves an L^2 bound.

Without the timelike assumption, a direct calculation in terms of the metric coefficients gives

$$\begin{aligned} \frac{2}{\sqrt{A}} \tilde{\mathbb{T}}(\partial_{t^*}, N_t) &= g^{t^*t^*} |Tv|^2 - g^{rr} |\tilde{\partial}_r v|^2 - g^{r\phi^*} \operatorname{Re}(\tilde{\partial}_r v \cdot \Phi \bar{v}) \\ &\quad - g^{\phi^*\phi^*} |\Phi v|^2 - g^{\theta\theta} |\partial_\theta v|^2 + (\nu^2 - n^2/4 + F) |v|^2, \end{aligned}$$

as well as

$$\frac{1}{\sqrt{A}} \tilde{\mathbb{T}}(\partial_{\phi^*}, N_t) = g^{t^*t^*} \operatorname{Re}(\Phi v \cdot T\bar{v}) + g^{t^*r} \operatorname{Re}(\Phi v \cdot \tilde{\partial}_r \bar{u}) + g^{t^*\phi^*} |\Phi v|^2.$$

If $v \in \mathcal{D}'_0(X_0)$, then $\Phi v = 0$ and hence

$$\tilde{\mathbb{T}}(K, N_t) = \frac{\sqrt{A}}{2} \left(g^{t^*t^*} |Tv|^2 - g^{rr} |\tilde{\partial}_r v|^2 - g^{\theta\theta} |\partial_\theta v|^2 + (\nu^2 - 9/4 + F) |v|^2 \right). \quad (4.1)$$

Although g^{rr} vanishes at $r = r_+$, the coefficients of $|\tilde{\partial}_r v|^2$ and $|\partial_\theta v|^2$ are certainly nonnegative on X_0 , while the coefficient of $|Tv|^2$ is strictly positive.

Proof of Proposition 2.1. If $v = e^{-i\lambda t^*} u$ for an axisymmetric function on u , then the contribution from $\tilde{\mathbb{T}}(K, N_t)$ will control $|\lambda|^2 \|u\|_{L^2(X_0)}^2$ for $|\lambda|$ large since the coefficient of $|Tv|^2$ is strictly positive. The proof can now be finished as in Lemma 3.9. $\square$

REFERENCES

- [1] Venkat Balasubramanian, Alex Buchel, Stephen R Green, Luis Lehner, and Steven L Liebling. Holographic thermalization, stability of anti-de sitter space, and the fermi-pasta-ulam paradox. *Physical review letters*, 113(7):071601, 2014.
- [2] P. Bizoń. Is AdS stable? *Gen.Rel.Grav.*, 46(5):1724, 2014.
- [3] P. Bizoń and A. Rostworowski. Weakly turbulent instability of anti de sitter spacetime. *Phys. Rev. Lett.*, 107:031102, Jul 2011.
- [4] Piotr Bizoń, Maciej Maliborski, and Andrzej Rostworowski. Resonant dynamics and the instability of anti-de sitter spacetime. *Physical review letters*, 115(8):081103, 2015.
- [5] Alex Buchel, Stephen R Green, Luis Lehner, and Steven L Liebling. Conserved quantities and dual turbulent cascades in anti-de sitter spacetime. *Physical Review D*, 91(6):064026, 2015.
- [6] Ben Craps, Oleg Evnin, and Joris Vanhoof. Renormalization group, secular term resummation and AdS (in)stability. *JHEP*, 10:048, 2014.
- [7] Ben Craps, Oleg Evnin, and Joris Vanhoof. Renormalization, averaging, conservation laws and AdS (in)stability. *JHEP*, 01:108, 2015.
- [8] Ó.J.C. Dias, G.T. Horowitz, D. Marolf, and J.E. Santos. On the nonlinear stability of asymptotically anti-de sitter solutions. *Class.Quant.Grav.*, 29:235019, 2012.
- [9] Ó.J.C Dias, G.T. Horowitz, and J.E. Santos. Gravitational turbulent instability of anti-de sitter space. *Classical and Quantum Gravity*, 29(19):194002, 2012.
- [10] Dominic Dold. Unstable mode solutions to the Klein-Gordon equation in Kerr-anti-de Sitter spacetimes. 2015.
- [11] S. Dyatlov and M. Zworski. Mathematical theory of scattering resonances.
- [12] G. Festuccia and H. Liu. A Bohr-Sommerfeld quantization formula for quasinormal frequencies of AdS black holes. *Advanced Science Letters*, 2(2):221–235, 2009.
- [13] O. Gannot. A global definition of quasinormal modes for Kerr-AdS Black Holes.
- [14] O. Gannot. Quasinormal modes for Schwarzschild-AdS black holes: Exponential convergence to the real axis. *Communications in Mathematical Physics*, 330(2):771–799, 2014.
- [15] O. Gannot. Elliptic boundary value problems for Bessel operators, with applications to anti-de Sitter spacetimes. July 2015. arXiv:1507.02794.
- [16] Israel Gohberg and Mark Grigorevich Kreĭn. *Introduction to the theory of linear nonselfadjoint operators*, volume 18. American Mathematical Soc., 1969.
- [17] G. Holzegel. Well-posedness for the massive wave equation on asymptotically anti-de sitter spacetimes. *Journal of Hyperbolic Differential Equations*, 09(02):239–261, June 2012.
- [18] G. Holzegel and J. Smulevici. Decay properties of klein-gordon fields on kerr-ads spacetimes. *Communications on Pure and Applied Mathematics*, 66(11):1751–1802, 2013.
- [19] G. Holzegel and J. Smulevici. Quasimodes and a lower bound on the uniform energy decay rate for kerr-ads spacetimes. *Analysis & PDE*, 7(5):1057–1090, 2014.
- [20] G.H. Holzegel and C.M. Warnick. Boundedness and growth for the massive wave equation on asymptotically anti-de Sitter black holes. 2012.
- [21] Gustav Holzegel. On the massive wave equation on slowly rotating kerr-ads spacetimes. *Communications in Mathematical Physics*, 294(1):169–197, 2010.
- [22] L. Hörmander. *The Analysis of Linear Partial Differential Operators III: Pseudo-Differential Operators*. A Series of Comprehensive Studies in Mathematics / Grundlehren Der Mathematischen Wissenschaften. Springer Berlin Heidelberg, 1985.
- [23] Alexandru D Ionescu and Sergiu Klainerman. On the uniqueness of smooth, stationary black holes in vacuum. *Inventiones mathematicae*, 175(1):35–102, 2009.

- [24] Boris I. Akovlevich Levin. *Distribution of zeros of entire functions*, volume 5. American Mathematical Soc., 1964.
- [25] R.B Melrose. Spectral and scattering theory for the laplacian on asymptotically euclidian spaces. *Lecture Notes in Pure and Applied Mathematics*, pages 85–85, 1994.
- [26] Johannes Sjostrand and Maciej Zworski. Complex scaling and the distribution of scattering poles. *Journal of the American Mathematical Society*, pages 729–769, 1991.
- [27] P. Stefanov. Quasimodes and resonances: sharp lower bounds. *Duke mathematical journal*, 99(1):75–92, 1999.
- [28] Plamen Stefanov. Approximating resonances with the complex absorbing potential method. *Communications in Partial Differential Equations*, 30(12):1843–1862, 2005.
- [29] S. Tang and M.j Zworski. From quasimodes to resonances. *Mathematical Research Letters*, 5:261–272, 1998.
- [30] A. Vasy. Microlocal analysis of asymptotically hyperbolic and kerr-de sitter spaces (with an appendix by semyon dyatlov). *Inventiones mathematicae*, 194(2):381–513, 2013.
- [31] C. M. Warnick. The Massive Wave Equation in Asymptotically AdS Spacetimes. *Communications in Mathematical Physics*, 321(1):85–111, May 2013.
- [32] C.M. Warnick. On quasinormal modes of asymptotically anti-de sitter black holes. *Communications in Mathematical Physics*, 333(2):959–1035, 2015.

E-mail address: ogannot@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, EVANS HALL, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA